

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" LABEL NUMBER: **EV 047297808 US**

DATE OF DEPOSIT: **January 28, 2002**

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE "EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER 37 CFR 1.10 ON THE DATE INDICATED ABOVE AND IS ADDRESSED TO:
THE COMMISSIONER OF PATENTS, BOX PATENT APPLICATION, WASHINGTON, D.C. 20231

JASON BERRY

(TYPED OR PRINTED NAME OF PERSON MAILING PAPER)


(SIGNATURE OF PERSON MAILING PAPER OR FEE)

**CONTINUATION-IN-PART
APPLICATION**

for

UNITED STATES LETTERS PATENT

on

**ABERRANTLY METHYLATED GENES AS
MARKERS OF BREAST MALIGNANCY**

by

**Saraswati Sukumar, Ella Evron, William C. Dooley,
Nicoletta Sacchi, Nancy Davidson and Mary Jo Fackler**

Sheets of Drawings: Twenty-Eight (28)
Docket No.: JHU1630-1

Attorneys

GRAY CARY WARE & FREIDENRICH LLP
4365 Executive Drive, Suite 1100
San Diego, California 92121-2133

10059579-042802

ABERRANTLY METHYLATED GENES AS MARKERS OF BREAST MALIGNANCY

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0001] This invention was made with Government support under P50 CA88843-01, awarded by the National Institutes of Health; and U.S. Public health Service Grant CA48943. The government may have certain rights in the invention.

RELATED APPLICATIONS

[0002] This is a continuation-in-part application of U.S. Patent Application Serial No. 09/771,337, filed January 26, 2001, co-pending, which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention relates generally to a method of diagnosing a cell proliferative disorder of breast tissue by determining the DNA methylation status of nucleic acids obtained a subject.

BACKGROUND

[0004] Methylation has been shown by several lines of evidence to play a role in gene activity, cell differentiation, tumorogenesis, X-chromosome inactivation, genomic imprinting and other major biological processes (Razin, A., H., and Riggs, R.D. eds. in DNA Methylation Biochemistry and Biological Significance, Springer-Verlag, New York, 1984). In eukaryotic cells, methylation of cytosine residues that are immediately 5' to a guanosine, occurs predominantly in cytosine-guanine (CG) poor regions (Bird, Nature, 321:209, 1986). In contrast, CpG islands remain unmethylated in normal cells, except during X-chromosome

10059579-012002

inactivation (Migeon, *et al.*, *supra*) and parental specific imprinting (Li, *et al.*, *Nature*, 366:362, 1993) where methylation of 5' regulatory regions can lead to transcriptional repression. De novo methylation of the Rb gene has been demonstrated in a small fraction of retinoblastomas (Sakai, *et al.*, *Am. J. Hum. Genet.*, 48:880, 1991), and recently, a more detailed analysis of the VHL gene showed aberrant methylation in a subset of sporadic renal cell carcinomas (Herman, *et al.*, *Proc. Natl. Acad. Sci., U.S.A.*, 91:9700, 1994). Expression of a tumor suppressor gene can also be abolished by de novo DNA methylation of a normally unmethylated CpG island (Issa, *et al.*, *Nature Genet.*, 7:536, 1994; Herman, *et al.*, *supra*; Merlo, *et al.*, *Nature Med.*, 1:686, 1995; Herman, *et al.*, *Cancer Res.*, 56:722, 1996; Graff, *et al.*, *Cancer Res.*, 55:5195, 1995; Herman, *et al.*, *Cancer Res.*, 55:4525, 1995).

[0005] Human cancer cells typically contain somatically altered nucleic acid, characterized by mutation, amplification, or deletion of critical genes. In addition, the nucleic acid from human cancer cells often displays somatic changes in DNA methylation (Fearon, *et al.*, *Cell*, 61:759, 1990; Jones, *et al.*, *Cancer Res.*, 46:461, 1986; Holliday, *Science*, 238:163, 1987; De Bustros, *et al.*, *Proc. Natl. Acad. Sci., USA*, 85:5693, 1988); Jones, *et al.*, *Adv. Cancer Res.*, 54:1, 1990; Baylin, *et al.*, *Cancer Cells*, 3:383, 1991; Makos, *et al.*, *Proc. Natl. Acad. Sci., USA*, 89:1929, 1992; Ohtani-Fujita, *et al.*, *Onco-gene*, 8:1063, 1993). However, the precise role of abnormal DNA methylation in human tumorigenesis has not been established.

[0006] Aberrant methylation of normally unmethylated CpG islands has been described as a frequent event in immortalized and transformed cells, and has been associated with transcriptional inactivation of defined tumor suppressor genes in human cancers. This molecular defect has also been described in association with various cancers. CpG islands are short sequences rich in the CpG dinucleotide and can be found in the 5' region of about half of all human genes. Methylation of cytosine within 5' CGIs is associated with loss of gene expression and has been seen in physiological conditions such as X chromosome inactivation and genomic imprinting. Aberrant methylation of CpG islands has been detected in genetic diseases such as the fragile-X syndrome, in aging cells and in neoplasia. About half of the tumor suppressor genes which have been shown to be mutated in the germline of

10059579-012303

patients with familial cancer syndromes have also been shown to be aberrantly methylated in some proportion of sporadic cancers, including Rb, VHL, p16, hMLH1, and BRCA1 (reviewed in Baylin, *et al.*, Adv. Cancer Res. 72:141-196 1998). Methylation of tumor suppressor genes in cancer is usually associated with (1) lack of gene transcription and (2) absence of coding region mutation. Thus CpG island methylation can serve as an alternative mechanism of gene inactivation in cancer.

[0007] Breast cancer is by far the most common form of cancer in women, and is the second leading cause of cancer death in humans. Despite many recent advances in diagnosing and treating breast cancer, the prevalence of this disease has been steadily rising at a rate of about 1% per year since 1940. Today, the likelihood that a woman living in North America will develop breast cancer during her lifetime is one in eight.

[0008] Breast cancer is often discovered at a stage that is advanced enough to severely limit therapeutic options and survival rates. Accordingly, more sensitive and reliable methods are needed to detect small (less than 2 cm diameter), early stage, *in situ* carcinomas of the breast. In addition to the problem of early detection, there remain serious problems in distinguishing between malignant and benign breast disease, in staging known breast cancers, and in differentiating between different types of breast cancers (*e.g.*, estrogen dependent versus non-estrogen dependent tumors). Recent efforts to develop improved methods for breast cancer detection have focused on cancer markers such as proteins that are uniquely expressed (*e.g.*, as a cell surface or secreted protein) by cancerous cells, or are expressed at measurably increased or decreased levels by cancerous cells compared to normal cells. Accordingly, the use of the methylation status of certain genes as a marker of cancer or cancerous conditions provides an additional weapon in early detection and prognosis of breast cancers.

[0009] Identification of the earliest genetic changes in cells associated with breast cancer is a major focus in molecular cancer research. Diagnostic approaches based on identification of these changes in specific genes are likely to allow implementation of early detection strategies and novel therapeutic approaches. Targeting these early changes might lead to more effective cancer treatment.

40059579.0125002

SUMMARY OF THE INVENTION

[0010] The present invention is based on the finding that several genes are newly identified as being differentially methylated in breast cancers. This seminal discovery is useful for breast cancer screening, risk-assessment, prognosis, disease identification, disease staging and identification of therapeutic targets. The identification of new genes that are methylated in breast cancer allows accurate and effective early diagnostic assays, methylation profiling using multiple genes; and identification of new targets for therapeutic intervention.

[0011] In a first embodiment, the invention provides method of diagnosing a cellular proliferative disorder of breast tissue in a subject comprising determining the state of methylation of one or more nucleic acids isolated from the subject. The state of methylation of one or more nucleic acids compared with the state of methylation of one or more nucleic acids from a subject not having the cellular proliferative disorder of breast tissue is indicative of a cell proliferative disorder in the subject. In one aspect of this embodiment, the state of methylation is hypermethylation. The invention provides a method of diagnosing a cellular proliferative disorder of breast tissue in a subject by detecting the state of methylation of one or more of the following nucleic acids: Twist, cyclin D2, WT1, NES-1, HOXA5 and combinations thereof. Also methylated are RAR β 2, 14.3.3 sigma, estrogen receptor, RASSF1A, HIN-1 and combinations thereof. In one aspect of the invention, nucleic acids are methylated in regulatory regions.

[0012] Invention methods can be used to diagnose disorders of the breast including breast cancers. In one aspect of the invention, disorders of the breast include ductal carcinoma *in situ*, lobular carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, intraductal carcinoma *in situ*, lobular carcinoma *in situ* and papillary carcinoma *in situ*.

[0013] Another embodiment of the invention provides a method of determining a predisposition to a cellular proliferative disorder of breast tissue in a subject. The method includes determining the state of methylation of one or more nucleic acids isolated from the subject, wherein the state of methylation of one or more nucleic acids compared with the state

10059579.012802

of methylation of the nucleic acid from a subject not having a predisposition to the cellular proliferative disorder of breast tissue is indicative of a cell proliferative disorder of breast tissue in the subject. The nucleic acids can be nucleic acids encoding Twist, cyclin D2, RAR β 2, HOXA5, WT1, 14.3.3 sigma, estrogen receptor, NES-1, RASSF1A, HIN-1 and combinations thereof.

[0014] Still another embodiment of the invention provides a method for detecting a cellular proliferative disorder of breast tissue in a subject. The method includes contacting a specimen containing at least one nucleic acid from the subject with an agent that provides a determination of the methylation state of at least one nucleic acid. The method further includes identifying the methylation states of at least one region of at least one nucleic acid, wherein the methylation state of the nucleic acid is different from the methylation state of the same region of nucleic acid in a subject not having the cellular proliferative disorder of breast tissue.

[0015] Yet a further embodiment of the invention provides a kit useful for the detection of a cellular proliferative disorder in a subject comprising carrier means compartmentalized to receive a sample therein; and one or more containers comprising a first container containing a reagent that modifies unmethylated cytosine and a second container containing primers for amplification of a CpG-containing nucleic acid. The primers hybridize with target polynucleotide sequence having the sequence of certain nucleic acids described herein.

SUMMARY OF THE FIGURES

[0016] **Figure 1A** shows the nucleotide sequence of the cyclin D2 promoter (SEQ ID NO:105). Regions highlighted indicate primer sequences. CG nucleotide pairs are shown capitalized and bolded. A highlighted box shows the location of an atg codon. **Figure 1B** shows nucleotide sequences for forward (F) and reverse (R) primer external and internal pairs used to detect methylated (M) and unmethylated (U) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0017] **Figures 2A and 2B** show the nucleotide sequence of the TWIST promoter (SEQ ID NO:106). Regions highlighted indicate primer sequences. CG nucleotide pairs are shown capitalized and bolded. A highlighted box shows the location of an atg codon. **Figure 2C** shows nucleotide sequences for forward (F) and reverse (R) external and internal primer pairs used to detect methylated (M) and unmethylated (U) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0018] **Figure 3A** shows the nucleotide sequence of the Retinoic Acid Receptor Beta (RAR β) promoter (SEQ ID NO:91). Regions highlighted indicate primer sequences. CG nucleotide pairs are shown capitalized and bolded. A highlighted box shows the location of an atg codon. **Figure 3B** shows nucleotide sequences for forward (F) and reverse (R) external and internal primer pairs used to detect methylated (M) and unmethylated (U) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0019] **Figure 4A** shows the nucleotide sequence of *Homo sapiens* serine protease-like protease (NES-1) mRNA. **Figure 4B** shows the nucleotide sequence of the NES-1 region (exon 3) analyzed. Regions highlighted indicate primer sequences. CG nucleotide pairs are shown capitalized and bolded. **Figure 4C** shows nucleotide sequences for forward (F) and reverse (R) primer pairs used to detect methylated (M) and unmethylated (U) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0020] **Figure 5A** shows the nucleotide sequence of HOXA5 promoter (3' to 5'). CG nucleotide pairs are shown capitalized and bolded. A highlighted box shows the location of a cat codon. **Figure 5B** shows the nucleotide sequence of the complementary region (5' to 3') analyzed (nucleotides -97 to -303). Regions highlighted indicate primer sequences. CG nucleotide pairs are shown capitalized and bolded. Highlighted box shows an atg codon. **Figure 5C** shows nucleotide sequences for forward (F) and reverse (R) primer pairs used to detect methylated (M) and unmethylated (U) nucleic acids. The base pair (BP) length of the primer pair product is also indicated. **Figure 5D** shows forward and reverse (sense and antisense) primers used for sequencing and expression of HOXA5.

1005937-01E001

[0021] Figure 6A to 6F show the nucleotide sequence of *Homo sapiens* 14.3.3 sigma protein promoter and gene, complete cds.

[0022] Figure 7A and 7B show the nucleotide sequence of *Homo sapiens* Wilms' tumor (WT1) gene promoter.

[0023] Figure 8A and 8B show the nucleotide sequence of *Homo. sapiens* estrogen receptor beta gene, promoter region and partial cds. CG nucleotide pairs are shown capitalized and bolded. Figure 8C shows nucleotide sequences of forward (F) and reverse (R) primer pairs used to detect methylated (M) and unmethylated (UM) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0024] Figure 9A shows the nucleotide sequence of human HIN-1 cDNA Regions highlighted indicate primer sequences. Figure 9B shows nucleotide sequences of forward and reverse external and internal primer pairs used to detect methylated and unmethylated nucleic acids. The base pair (bp) length of the primer pair product is also indicated.

[0025] Figure 10A shows the nucleotide sequence of the RASSF1A promoter. CG nucleotide pairs are shown capitalized and bolded. Regions highlighted indicate primer sequences. Figure 10B shows nucleotide sequences of forward (F) and reverse (R) external and internal primer pairs used to detect methylated (M) and unmethylated (UM) nucleic acids. The base pair (BP) length of the primer pair product is also indicated.

[0026] Figure 11 is a schematic representation of the invention assay methods utilizing the technique of multiplex methylation-specific PCR.

DETAILED DESCRIPTION OF THE INVENTION

[0027] The invention is based upon the discovery that the hypermethylation of certain genes can serve as markers for cellular proliferative disorders of breast tissue. This is the first time that promoter hypermethylation of certain genes such as Twist, cyclin D2, RAR β 2, WT1, NES-1 and HOXA5 have been associated with breast cancer.

10059579 012002

[0028] It has been determined that the methylation state of nucleic acids of certain genes, particularly regulatory sequences, is diagnostic for the presence or potential development of a cellular proliferative disorder of breast tissue in subjects. More particularly, the hypermethylation of certain nucleotides localized in CpG islands has been shown to affect the expression of genes associated with the CpG islands; typically such hypermethylated genes have reduced or abolished expression, primarily due to down-regulated transcription. Hypermethylation of, for example, Twist, cyclin D2, retinoic acid receptor β (RAR β), WT1, HOXA5, 14.3.3 sigma, estrogen receptor (ER) NES-1, the Ras association domain family 1A gene (RASSF1A), and the high in normal-1 gene (HIN-1) allows one to diagnose a cellular proliferative disorder of breast tissue. Using a recently developed PCR-based technique called methylated specific PCR (MSP), aberrantly methylated nucleic acids in breast cancer primary tumors and biological samples from individuals with breast cancer can be identified.

[0029] In a first embodiment, the invention provides a method of diagnosing a cellular proliferative disorder of breast tissue in a subject comprising determining the state of methylation of one or more nucleic acids isolated from the subject, wherein the state of methylation of one or more nucleic acids as compared with the state of methylation of one or more nucleic acids from a subject not having the cellular proliferative disorder of breast tissue is indicative of a cellular proliferative disorder of breast tissue in the subject. A preferred nucleic acid is a CpG-containing nucleic acid, such as a CpG island.

[0030] A cell proliferative disorder as described herein may be a neoplasm. Such neoplasms are either benign or malignant. The term "neoplasm" refers to a new, abnormal growth of cells or a growth of abnormal cells that reproduce faster than normal. A neoplasm creates an unstructured mass (a tumor), which can be either benign or malignant. The term "benign" refers to a tumor that is noncancerous, e.g. its cells do not invade surrounding tissues or metastasize to distant sites. The term "malignant" refers to a tumor that is metastatic, invades contiguous tissue or no longer under normal cellular growth control.

[0031] One type of cellular proliferative disorder is a cell proliferative disorder of breast tissue. Disorders of breast tissue or breast cancers can involve numerous cells and tissues resulting in various disorders of the breast including ductal carcinoma *in situ*, lobular

10059579.012302

carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, intraductal carcinoma *in situ*, lobular carcinoma *in situ*, and papillary carcinoma *in situ*.

[0032] The invention method includes determining the state of methylation of one or more nucleic acids isolated from the subject. The phrases "nucleic acid" or "nucleic acid sequence" as used herein refer to an oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent a sense or antisense strand, peptide nucleic acid (PNA), or to any DNA-like or RNA-like material, natural or synthetic in origin. As will be understood by those of skill in the art, when the nucleic acid is RNA, the deoxynucleotides A, G, C, and T are replaced by ribonucleotides A, G, C, and U, respectively.

[0033] The nucleic acid of interest can be any nucleic acid where it is desirable to detect the presence of a differentially methylated CpG island. The CpG island is a CpG rich region of a nucleic acid sequence. The nucleic acids includes, for example, a sequence encoding the following genes (GenBank Accession Numbers are shown, followed by the nucleotides corresponding to the region(s) examined for the presence or absence of methylation (numbers are relative to the first ATG codon unless otherwise indicated)): Twist (Accession No. AC003986; -51145 to 151750 (complement) (SEQ ID NO:106) , cyclin D2 (Accession No. U47284; -1616 to -1394) (SEQ ID NO:105); RAR β 2 (Accession No. AF; 157484; -196 to -357)(SEQ ID NO:91), WT1 (Accession No. AB034940) (SEQ ID NO:103); HOXA5 (Accession No. AC004080) (SEQ ID NO:96), 14.3.3 sigma (Accession No. AF029081) (SEQ ID NO:102); estrogen receptor (ER; Accession No. X62462) (SEQ ID NO:104); NES-1 (Accession No. AF024605) (SEQ ID NO:94); RASSF1A (Accession No. AF102770) (SEQ ID NO:121); and HIN-1 (Accession No. AY040564) (SEQ ID NO:120), the nucleotide sequence of each of which is incorporated by reference herein.

[0034] WT1 encodes a transcriptional regulatory protein that binds DNA via four Cys₂.His₂ zinc fingers. WT1 mRNA undergoes two independent splicing events leading to the expression of at least four predominant isoforms. These splices result in the inclusion or

10059579.01E306

omission of exon 5 (51 base pairs) and the presence or absence of a nine base pair insert (encoding three amino acids, KTS) between the third and fourth zinc finger domains. Lack of expression has been observed in some Wilms' tumors, leading to classification as a tumor suppressor gene. However, WT1 is overexpressed in 75% of cases of acute leukemia and is upregulated as chronic myeloid leukemia progresses into blast crisis. Thus, WT1 can apparently be either a tumor suppressor or an oncogene.

[0035] The cyclin D1, D2 and D3 proteins are involved in regulation of the cell cycle through phosphorylation and inactivation of the retinoblastoma protein and activation of cyclin E, leading to transition of the cells from G1 to DNA synthesis. In addition to their role in cell cycle regulation, the D-type cyclins have been implicated in differentiation and neoplastic transformation. Overexpression of cyclin D2 has been reported in gastric cancer, and was shown to correlate with disease progression and poor prognosis. Overexpression of cyclin D2 is also noted in ovarian granulosa cell tumors and testicular germ cell tumor cell lines.

[0036] 14.3.3 σ is a member of a superfamily that is responsible for instituting the G2 cell cycle checkpoint in response to DNA damage in human cells (Hermeking, *et al.* (1997) *Mol. Cell* 1, 3-11; Chan, *et al.* (1999) *Nature* 401, 616-20). In addition to any growth advantage resulting from a loosening of this checkpoint control mechanism, loss of σ function is predicted to cause an increase in DNA damage in response to γ -irradiation. Loss of 14.3.3. σ in primary epithelial cells leads to immortalization (Dellambra *et al.* (2000) *J. Cell Biol.*, 149:1117-1130), one of the earliest steps towards cancer.

[0037] Retinoic acid (RA) controls fundamental developmental processes, induces terminal differentiation of myeloid progenitors and suppresses cancer and cell growth. RA activity is mediated by nuclear receptors, the retinoic acid receptors, RARs, that act as RA-dependent transcriptional activators in their heterodimeric forms with retinoid X receptors, RXRs (Chambon, 1996). RARs induce local chromatin changes at level of target genes, containing responsive RA elements (RAREs) by recruiting multiprotein complexes with histone acetyltransferase (HAT) activity and histone deacetylase (HDAC) activity that dynamically pattern chromatin modification and regulate gene expression. RARs and RXRs,

1005579-012802
200210-1625001

when disrupted, result in severe developmental defects and neoplastic transformation. In breast cancer cells, the expression of one member of the RARs family, RAR β is found consistently down regulated or lost. RAR β downregulation can be reversed by RA in estrogen receptor (ER)-positive, but not in ER-negative breast carcinoma cell lines, believed to represent more advanced forms of tumors. Loss of RA-induced RAR β expression is considered a crucial step in the development of RA-resistance in breast carcinogenesis. A complex regulatory region, with two promoters, regulates RAR β gene expression. Only one promoter, RAR β 2, containing several RA-response elements, including a canonical and an auxiliary RA response element, β RARE, is active in human mammary epithelial cells (HMEC). The transcription of the RAR β 2 promoter is mediated by multiple RARs including RAR α and RAR β itself able to recruit coactivator and corepressor protein complexes with HAT/HDAC activities, respectively.

[0038] The Twist gene product is a transcription factor with DNA binding and helix-loop-helix domains. Twist is a member of the bHLH transcription factor family and is involved in the development of mesodermally derived tissue including the skeleton. In humans, mutations in the Twist gene have been identified in patients with Saethre-chotzen syndrome, a relatively common craniosynostosis disorder with autosomal dominant inheritance. (see Gripp *et al.*, (2000) *Hum. Mutat.* 15:479.) Twist also influences osteogenic gene expression and may act as a master switch in initiating bone cell differentiation by regulating the osteogenic cell lineage (Lee *et al.*, (1999) *J. Cell Biochem.* 75:566-577).

[0039] NES1 (normal epithelial cell-specific 1) is a novel gene with a predicted polypeptide of about 30.14 kilodaltons and having a 50-63% similarity and 34-42% identity with several families of serine proteases, in particular the trypsin-like proteases, members of the glandular kallikrein family (including prostate-specific antigen, nerve growth factor gamma, and epidermal growth factor-binding protein) and the activators for the kringle family proteins (including the human tissue plasminogen activator and human hepatocyte growth factor activator) (Liu *et al.*, (1996) *Cancer Res.* 56:14 3371-9). All of the residues known to be crucial for substrate binding, specificity, and catalysis by the serine proteases are conserved in the predicted NES1 protein, indicating that it has protease-like activity.

10059579.012802

Immunolocalization studies with an antipeptide antibody directed against a unique region of the NES1 protein (amino acids 120-137) detect a specific 30-kilodalton polypeptide almost exclusively in the supernatant of the mRNA-positive mammary epithelial cells (MECs), suggesting that NES1 is a secreted protein. The 1.4-kb NES1 mRNA is expressed in several organs (thymus, prostate, testis, ovary, small intestine, colon, heart, lung, and pancreas) with highest levels in the ovaries. Although expression of the NES1 mRNA is observed in all normal and immortalized nontumorigenic MECs, the majority of human breast cancer cell lines show a drastic reduction or a complete lack of its expression. The structural similarity of NES1 to polypeptides known to regulate growth factor activity and a negative correlation of NES1 expression with breast oncogenesis suggest a direct or indirect role for this novel protease-like gene product in the suppression of tumorigenesis. Studies using fluorescence in situ hybridization localized the NES1 gene to chromosome 19q13.3, a region that contains genes for related proteases (Goyal *et al.*, (1998) *Cancer Res.*, 58:21 4782-6).

[0040] The HOX genes are expressed during embryonic development and have a role in specifying antero-posterior positional information. The genes are arranged in four clusters and a collinear relation exists between a gene's position in the cluster and its anterior boundary of expression. Genes with more anterior boundaries are also expressed earlier than genes with more posterior boundaries. Hox genes encode transcription factors; therefore, a model for the coordinate regulation of the genes within the HOX clusters is that Hox gene products regulate their own expression. The production of HOXA5 from an expression vector can activate a transient and simultaneous expression of other upstream and downstream genes of the same HOX cluster and genes from other clusters.

[0041] The estrogen receptor gene has been implicated in the initiation and/or progression of human breast cancer. Loss of expression of either gene has been associated with poorly differentiated tumors and poorer prognosis. Several studies have reported an association between estrogen receptor (ER) expression and breast tumors. A loss of ER expression has been associated with aberrant 5' CpG island methylation in breast cancer cell lines and primary human breast tumors. Studies show that aberrant methylation of ER CpG islands begins before invasion of tumors into surrounding tissues and it increases with metastatic

progression (Naas *et al.*, (2000) *Cancer Res.*, 60:4346-4348; incorporated by reference in its entirety).

[0042] Hypermethylation of the CpG island of Ras Association Domain Family 1A (RASSF1A), a putative tumor suppressor gene from the 3p21.3 locus, occurs in a large percentage of human breast cancers. Hypermethylation of the RASSF1A promoter appears to be the main mechanism of inactivation. The high frequency of epigenetic inactivation of the RASF1A gene in breast cancer supports its role as a putative tumor suppressor gene (R. Dammann, *et al.*, *Cancer Research* 61:3105-3109, 2001; K Dreijerink *et al.*, *PNAS* 98(18):7504-7509, 2001; D.G. Burbee *et al.*, *J. National Cancer Institute* 93(9):691-699, 2001, each of which is incorporated herein by reference in its entirety).

[0043] Expression of HIN-1 (high in normal-1) is significantly down regulated in 94% of human breast carcinoma and in 95% of preinvasive lesions, such as ductal and lobular carcinoma in situ. This decrease in HIN-1 expression is accompanied by hypermethylation of its promoter in the majority of breast cancer cell lines and primary tumors. This decrease in HIN-1 expression is accompanied by hypermethylation of its promoter in the majority of breast cancer cell lines (greater than 90%) and primary tumors (74%). HIN-1 is a putative cytokine with no significant homology to known proteins. Reintroduction of HIN-1 into breast cancer cells has been shown to inhibit cell growth, making HIN-1 a candidate tumor suppressor gene that is inactivated at high frequency in the earliest stages of breast tumorigenesis (I.E. Krop *et al.*, *PNAS* 98(17):9796-9801, 2001, which is incorporated herein by reference in its entirety).

[0044] Any nucleic acid sample, in purified or nonpurified form, can be utilized in accordance with the present invention, provided it contains, or is suspected of containing, a nucleic acid sequence containing a target locus (e.g., CpG-containing nucleic acid). One nucleic acid region capable of being differentially methylated is a CpG island, a sequence of nucleic acid with an increased density relative to other nucleic acid regions of the dinucleotide CpG. The CpG doublet occurs in vertebrate DNA at only about 20% of the frequency that would be expected from the proportion of G•C base pairs. In certain regions, the density of CpG doublets reaches the predicted value; it is increased by ten fold relative to

100597-010001
2002-06-25

the rest of the genome. CpG islands have an average G•C content of about 60%, compared with the 40% average in bulk DNA. The islands take the form of stretches of DNA typically about one to two kilobases long. There are about 45,000 such islands in the human genome.

[0045] In many genes, the CpG islands begin just upstream of a promoter and extend downstream into the transcribed region. Methylation of a CpG island at a promoter usually prevents expression of the gene. The islands can also surround the 5' region of the coding region of the gene as well as the 3' region of the coding region. Thus, CpG islands can be found in multiple regions of a nucleic acid sequence including upstream of coding sequences in a regulatory region including a promoter region, in the coding regions (*e.g.*, exons), downstream of coding regions in, for example, enhancer regions, and in introns.

[0046] In general, the CpG-containing nucleic acid is DNA. However, invention methods may employ, for example, samples that contain DNA, or DNA and RNA, including messenger RNA, wherein DNA or RNA may be single stranded or double stranded, or a DNA-RNA hybrid may be included in the sample. A mixture of nucleic acids may also be employed. The specific nucleic acid sequence to be detected may be a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid. It is not necessary that the sequence to be studied be present initially in a pure form; the nucleic acid may be a minor fraction of a complex mixture, such as contained in whole human DNA. The nucleic acid-containing sample used for determination of the state of methylation of nucleic acids contained in the sample or detection of methylated CpG islands may be extracted by a variety of techniques such as that described by Sambrook, *et al.* (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, NY, 1989; incorporated in its entirety herein by reference).

[0047] A nucleic acid can contain a regulatory region which is a region of DNA that encodes information that directs or controls transcription of the nucleic acid. Regulatory regions include at least one promoter. A "promoter" is a minimal sequence sufficient to direct transcription, to render promoter-dependent gene expression controllable for cell-type specific, tissue-specific, or inducible by external signals or agents. Promoters may be located

1005579.012802

in the 5' or 3' regions of the gene. Promoter regions, in whole or in part, of a number of nucleic acids can be examined for sites of CG-island methylation.

[0048] Nucleic acids isolated from a subject are obtained in a biological specimen from the subject. The nucleic acid may be isolated from breast tissue, blood, plasma serum, lymph, duct cells, nipple aspiration fluid, ductal lavage fluid and bone marrow. Tissue, blood, lymph, lymph node, duct cells, nipple aspiration fluid, ductal lavage fluid and bone marrow are obtained by various medical procedures known to those of skill in the art. Duct cells can be obtained by nipple aspiration, ductal lavage, sentinel node biopsy, fine needle aspirate, routine operative breast endoscopy and core biopsy. Ductal lavage fluid can be obtained by using a DucWash procedure. In this procedure, a catheter is inserted into one or more of the four to eight ducts typically present in each human breast, lavage of the duct is performed, and the lavage fluid is collected. Alternatively, ductal lavage may be achieved through a microcatheter procedure known as ROBE (routine operative breast endoscopy), which allows visualization of a tumor at the same time as aspiration of fluid from the duct.

[0049] In one aspect of the invention, the state of methylation in nucleic acids of the sample obtained from a subject is hypermethylation compared with the same regions of the nucleic acid in a subject not having the cellular proliferative disorder of breast tissue. Hypermethylation, as used herein, is the presence of methylated alleles in one or more nucleic acids. Nucleic acids from a subject not having a cellular proliferative disorder of breast tissues contain no detectable methylated alleles when the same nucleic acids are examined.

[0050] A method for determining the methylation state of nucleic acids is described in U.S. Patent No. 6,017,704 which is incorporated herein in its entirety and described briefly herein. Determining the methylation state of the nucleic acid includes amplifying the nucleic acid by means of oligonucleotide primers that distinguishes between methylated and unmethylated nucleic acids.

[0051] Two or more markers can also be screened simultaneously in a single amplification reaction to generate a low cost, reliable cancer-screening test for breast cancers. A

2025-05-01 10:55:00

combination of DNA markers for CpG-rich regions of nucleic acid may be amplified in a single amplification reaction. The markers are multiplexed in a single amplification reaction, for example, by combining primers for more than one locus. For example, DNA from a ductal lavage sample can be amplified with two or more different unlabeled or randomly labeled primer sets in the same amplification reaction. Especially useful are two or more markers selected from cyclin D2, RAR β 2, Twist, NES-1, RASSF1A and HIN-1. The reaction products are separated on a denaturing polyacrylamide gel, for example, and then exposed to film or stained with ethidium bromide for visualization and analysis. By analyzing a panel of markers, there is a greater probability of producing a more useful methylation profile for a subject.

[0052] For example, a screening technique, referred to herein as "multiplex methylation-specific PCR" is a unique version of methylation-specific PCR. Methylation-specific PCR is described in U.S. Patent Nos. 5,786,146, 6,200,756, 6,017,704 and 6,265,171, each of which is incorporated herein by reference in its entirety. Multiplex methylation-specific PCR utilizes MSP primers for a multiplicity of markers, for example up to five different breast cancer markers, in a two-stage nested PCR amplification reaction. The primers used in the first PCR reaction are selected to amplify a larger portion of the target sequence than the primers of the second PCR reaction. The primers used in the first PCR reaction are referred to herein as "external primers" or DNA primers" and the primers used in the second PCR reaction are referred to herein as "MSP primers." Two sets of primers (i.e., methylated and unmethylated for each of the markers targeted in the reaction) are used as the MSP primers. In addition in multiplex methylation-specific PCR, as described herein, a small amount (i.e., 1 μ l) of a 1:10¹ to about 10⁶ dilution of the reaction product of the first "external" PCR reaction is used in the second "internal" MSP PCR reaction. The technique of multiplex methylation-specific PCR is illustrated schematically in Figure 11.

[0053] As shown in Table 1 below, multiplex methylation-specific PCR greatly enhances the accuracy of diagnosis obtainable from an amount of DNA available for analysis as compared with direct PCR analysis.

10050570-010000

Table 1

Method of PCR	DNA Sample	Usage calculations	Test Capacity
DIRECT MSP:	20 μ l DNA ($\leq 1 \mu$ g)	1 μ l per PCR rxn. 2 μ l per test. Sufficient for 20 rxns (10 tests); 2 replicate tests of 5 genes.	If all 20 μ l DNA sample is used, 10 tests evaluate 5 genes X 2
MULTIPLEX MSP:	20 μ l DNA ($\leq 1 \mu$ g)	2 μ l per 1 st PCR rxn (25 μ l PCR rxn). 1 μ l 10 ¹ dilution into 2 nd PCR rxn ($\leq 1 \mu$ g)	If 2 μ l DNA sample is used, 125 tests evaluate 5 genes X 25.
	If all 20 μ l starting DNA is used in multiplex methylation-specific PCR, up to 10 panels of 5 genes X 25 replicates. 2 μ l starting DNA is sufficient for 250 2 nd PCR rxns (0.1 μ l/rxn, 2 rxn/test, 125 tests from 25 μ l 1 st rxn		

[0054] Multiplex methylation-specific PCR is also high specific. Tests conducted to compare the results of direct MSP with multiplex methylation-specific PCR in analysis of the methylation status of human primary breast tumor, and human breast cancer cell lines, are summarized, respectively, in Tables 5-7 below. The results shown in Tables 5-7 illustrate concordance in the results obtained by analysis of these various types of samples using direct MPC and multiplex methylation-specific PCR, as disclosed herein.

[0055] If the sample is impure (e.g., plasma, serum, lymph, ductal cells, nipple aspiration fluid, ductal lavage fluid, bone marrow, blood or breast tissue embedded in paraffin), it may be treated before amplification with a reagent effective for lysing the cells contained in the fluids, tissues, or animal cell membranes of the sample, and for exposing the nucleic acid(s) contained therein. Methods for purifying or partially purifying nucleic acid from a sample are well known in the art (e.g., Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 1989, herein incorporated by reference).

[0056] Primers hybridize with target polynucleotide sequences. Nucleic acid sequences including exemplary primers are set forth in SEQ ID NO:1 to SEQ ID NO:128. Oligonucleotide primers specifically targeted to methylated and unmethylated genes including Twist, cyclinD2, RAR β 2, WT1, HOXA5, 14.3.3 sigma, estrogen receptor, NES-1, RASSF1A, HIN-1, and their associated CpG islands include, respectively, SEQ ID NO:7-14, 21-24, 37-40, 49-64, 69-72, 77-80, 85-90, 107-110, 116-119, 124-128, 129-130, and 135-136. (See Table 4 below).

[0057] Detection of differential methylation can be accomplished by contacting a nucleic acid sample with a methylation sensitive restriction endonuclease that cleaves only unmethylated CpG sites under conditions and for a time to allow cleavage of unmethylated nucleic acid. The sample is further contacted with an isoschizomer of the methylation sensitive restriction endonuclease that cleaves both methylated and unmethylated CpG-sites under conditions and for a time to allow cleavage of methylated nucleic acid. Oligonucleotides are added to the nucleic acid sample under conditions and for a time to allow ligation of the oligonucleotides to nucleic acid cleaved by the restriction endonuclease, and the digested nucleic acid is amplified by conventional methods, such as PCR wherein primers complementary to the oligonucleotides are employed. Following identification, the methylated CpG-containing nucleic acid can be cloned, using methods well known to those of skill in the art (see Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 1989).

[0058] As used herein, a "methylation sensitive restriction endonuclease" is a restriction endonuclease that includes CG as part of its recognition site and has altered activity when the C is methylated as compared to when the C is not methylated. Preferably, the methylation sensitive restriction endonuclease has inhibited activity when the C is methylated (e.g., SmaI). Specific non-limiting examples of methylation sensitive restriction endonucleases include Sma I, BssHII, or HpaII, MspI, BSTUI, and NotI. Such enzymes can be used alone or in combination. Other methylation sensitive restriction endonucleases will be known to those of skill in the art and include, but are not limited to SacII, and EagI, for example. An "isoschizomer" of a methylation sensitive restriction endonuclease is a restriction

1005579.012003

endonuclease that recognizes the same recognition site as a methylation sensitive restriction endonuclease but cleaves both methylated and unmethylated CGs. Those of skill in the art can readily determine appropriate conditions for a restriction endonuclease to cleave a nucleic acid (see Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, 1989).

[0059] A nucleic acid of interest is cleaved with a methylation sensitive endonuclease. Cleavage with the methylation sensitive endonuclease creates a sufficient overhang on the nucleic acid of interest, i.e., sufficient to allow specific hybridization of an oligonucleotide of interest. Following cleavage with the isoschizomer, the cleavage product can still have a sufficient overhang. An "overhang" refers to nucleic acid having two strands wherein the strands end in such a manner that a few bases of one strand are not base paired to the other strand. A "sufficient overhang" refers to an overhang of at least two bases in length or four or more bases in length. An overhang of a specific sequence on the nucleic acid of interest may be desired in order for an oligonucleotide of interest to hybridize. In this case, the isoschizomer can be used to create the overhang having the desired sequence on the nucleic acid of interest.

[0060] Cleavage with a methylation sensitive endonuclease results in a reaction product of the nucleic acid of interest that has a blunt end or an insufficient overhang. "Blunt end" refers to a flush ending of two stands, the sense stand and the antisense strand, of a nucleic acid. Once a sufficient overhang is created on the nucleic acid of interest, an oligonucleotide is ligated to the nucleic acid of interest, which has been cleaved by the methylation specific restriction endonuclease. "Ligation" is the attachment of two nucleic acid sequences by base pairing of substantially complementary sequences and/or by the formation of covalent bonds between two nucleic acid sequences.

[0061] An adaptor can be utilized to create DNA ends of desired sequence and overhang. An "adaptor" is a double-stranded nucleic acid sequence with one end that has a sufficient single-stranded overhang at one or both ends such that the adaptor can be ligated by base-pairing to a sufficient overhang on a nucleic acid of interest that has been cleaved by a methylation sensitive restriction enzyme or an isoschizomer of a methylation sensitive

200210645500T

restriction enzyme. Adaptors can be obtained commercially. Alternatively, two oligonucleotides that are substantially complementary over their entire sequence except for the region(s) at the 5' and/or 3' ends that will form a single stranded overhang can be used to form an adaptor. The single stranded overhang on the adaptor is selected to be complementary to an overhang on the nucleic acid cleaved by a methylation sensitive restriction enzyme or an isoschizomer of a methylation sensitive restriction enzyme, such that the overhang on the nucleic acid of interest will base pair with the 3' or 5' single stranded end of the adaptor under appropriate conditions. The conditions will vary depending on the sequence composition (GC vs AT), the length, and the type of nucleic acid (see Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed.; Cold Spring Harbor Laboratory Press, Plainview, NY, 1998).

[0062] Following the ligation of the oligonucleotide to the nucleic acid of interest, the nucleic acid of interest is amplified using a primer complementary to the oligonucleotide. Specifically, the term "primer" as used herein refers to a sequence comprising two or more deoxyribo-nucleotides or ribonucleotides, preferably more than three, and more preferably more than eight, wherein the sequence is capable of initiating synthesis of a primer extension product that is substantially complementary to a nucleic acid such as an adaptor or a ligated oligonucleotide. Environmental conditions conducive to synthesis include the presence of nucleoside triphosphates, an agent for polymerization, such as DNA polymerase, and suitable temperature and pH. The primer is preferably single stranded for maximum efficiency in amplification, but may be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. The primer can be an oligodeoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the agent for polymerization. The exact length of the primer will depend on many factors, including temperature, buffer composition (i.e., salt concentration), and nucleotide composition. The oligonucleotide primer typically contains 12-20 or more nucleotides, although it may contain fewer nucleotides.

[0063] Primers of the invention are designed to be "substantially" complementary to each strand of the oligonucleotide to be amplified and include the appropriate G or C nucleotides

as discussed above. This means that the primers must be sufficiently complementary to hybridize with their respective strands under conditions that allow the agent for polymerization to perform. In other words, the primers should have sufficient complementarity with a 5' and 3' oligonucleotide to hybridize therewith and permit amplification of CpG containing nucleic acid sequence.

[0064] Primers of the invention are employed in the amplification process, which is an enzymatic chain reaction that produces exponentially increasing quantities of target locus relative to the number of reaction steps involved (e.g., polymerase chain reaction or PCR). Typically, one primer is complementary to the negative (-) strand of the locus (antisense primer) and the other is complementary to the positive (+) strand (sense primer). Annealing the primers to denatured nucleic acid followed by extension with an enzyme, such as the large fragment of DNA Polymerase I (Klenow) and nucleotides, results in newly synthesized + and - strands containing the target locus sequence. Because these newly synthesized sequences are also templates, repeated cycles of denaturing, primer annealing, and extension results in exponential production of the region (i.e., the target locus sequence) defined by the primer. The product of the chain reaction is a discrete nucleic acid duplex with termini corresponding to the ends of the specific primers employed.

[0065] The oligonucleotide primers used in invention methods may be prepared using any suitable method, such as conventional phosphotriester and phosphodiester methods or automated embodiments thereof. In one such automated embodiment, diethylphosphoramidites are used as starting materials and may be synthesized as described by Beaucage, *et al.* (Tetrahedron Letters, 22:1859-1862, 1981). One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Patent No. 4,458,066.

[0066] Another method for detecting a methylated CpG-containing nucleic acid includes contacting a nucleic acid-containing specimen with an agent that modifies unmethylated cytosine, amplifying the CpG-containing nucleic acid in the specimen by means of CpG-specific oligonucleotide primers, wherein the oligonucleotide primers distinguish between modified methylated and non-methylated nucleic acid and detecting the methylated nucleic acid. The amplification step is optional and although desirable, is not essential. The

1005579-012302

method relies on the PCR reaction itself to distinguish between modified (e.g., chemically modified) methylated and unmethylated DNA.

[0067] The term "modifies" as used herein means the conversion of an unmethylated cytosine to another nucleotide that will facilitate methods to distinguish the unmethylated from the methylated cytosine. Preferably, the agent modifies unmethylated cytosine to uracil. Preferably, the agent used for modifying unmethylated cytosine is sodium bisulfite; however, other agents that similarly modify unmethylated cytosine, but not methylated cytosine, can also be used in the method. Sodium bisulfite (NaHSO_3) reacts readily with the 5,6-double bond of cytosine, but poorly with methylated cytosine. Cytosine reacts with the bisulfite ion to form a sulfonated cytosine reaction intermediate that is susceptible to deamination, giving rise to a sulfonated uracil. The sulfonate group can be removed under alkaline conditions, resulting in the formation of uracil. Uracil is recognized as a thymine by Taq polymerase. Therefore after PCR, the resultant product contains cytosine only at the position where 5-methylcytosine occurs in the starting template DNA.

[0068] The primers used in the invention for amplification of the CpG-containing nucleic acid in the specimen, after bisulfite modification, specifically distinguish between untreated or unmodified DNA, methylated, and non-methylated DNA. MSP primers for the non-methylated DNA preferably have a T in the 3' CG pair to distinguish it from the C retained in methylated DNA, and the complement is designed for the antisense primer. MSP primers usually contain relatively few Cs or Gs in the sequence since the Cs will be absent in the sense primer and the Gs absent in the antisense primer (C becomes modified to U (uracil) which is amplified as T (thymidine) in the amplification product).

[0069] The primers of the invention embrace oligonucleotides of sufficient length and appropriate sequence so as to provide specific initiation of polymerization on a significant number of nucleic acids in the polymorphic locus. Where the nucleic acid sequence of interest contains two strands, it is necessary to separate the strands of the nucleic acid before it can be used as a template for the amplification process. Strand separation can be effected either as a separate step or simultaneously with the synthesis of the primer extension products. This strand separation can be accomplished using various suitable denaturing

1005579.01E002

conditions, including physical, chemical, or enzymatic means, the word "denaturing" includes all such means. One physical method of separating nucleic acid strands involves heating the nucleic acid until it is denatured. Typical heat denaturation may involve temperatures ranging from about 80° to 105°C for times ranging from about 1 to 10 minutes. Strand separation may also be induced by an enzyme from the class of enzymes known as helicases or by the enzyme RecA, which has helicase activity, and in the presence of riboATP, is known to denature DNA. The reaction conditions suitable for strand separation of nucleic acids with helicases are described by Kuhn Hoffmann-Berling (CSH-Quantitative Biology, 43:63, 1978) and techniques for using RecA are reviewed in C. Radding (Ann. Rev. Genetics, 16:405-437, 1982).

[0070] When complementary strands of nucleic acids are separated, regardless of whether the nucleic acid was originally double or single stranded, the separated strands are ready to be used as a template for the synthesis of additional nucleic acid strands. This synthesis is performed under conditions allowing hybridization of primers to templates to occur. Generally synthesis occurs in a buffered aqueous solution, generally at a pH of about 7-9. Preferably, a molar excess (for genomic nucleic acid, usually about 108:1 primer:template) of the two oligonucleotide primers is added to the buffer containing the separated template strands. It is understood, however, that the amount of complementary strand may not be known if the process of the invention is used for diagnostic applications, so that the amount of primer relative to the amount of complementary strand cannot be determined with certainty. As a practical matter, however, the amount of primer added will generally be in molar excess over the amount of complementary strand (template) when the sequence to be amplified is contained in a mixture of complicated long-chain nucleic acid strands. Large molar excess is preferred to improve the efficiency of the process.

[0071] The deoxyribonucleoside triphosphates dATP, dCTP, dGTP, and dTTP are added to the synthesis mixture, either separately or together with the primers, in adequate amounts and the resulting solution is heated to about 90°-100°C from about 1 to 10 minutes, preferably from 1 to 4 minutes. After this heating period, the solution is allowed to cool to approximately room temperature, which is preferable for the primer hybridization. To the

100557-0100
200557-0100
300557-0100
400557-0100
500557-0100
600557-0100
700557-0100
800557-0100
900557-0100
100557-0100
110557-0100
120557-0100
130557-0100
140557-0100
150557-0100
160557-0100
170557-0100
180557-0100
190557-0100
200557-0100
210557-0100
220557-0100
230557-0100
240557-0100
250557-0100
260557-0100
270557-0100
280557-0100
290557-0100
300557-0100
310557-0100
320557-0100
330557-0100
340557-0100
350557-0100
360557-0100
370557-0100
380557-0100
390557-0100
400557-0100
410557-0100
420557-0100
430557-0100
440557-0100
450557-0100
460557-0100
470557-0100
480557-0100
490557-0100
500557-0100
510557-0100
520557-0100
530557-0100
540557-0100
550557-0100
560557-0100
570557-0100
580557-0100
590557-0100
600557-0100
610557-0100
620557-0100
630557-0100
640557-0100
650557-0100
660557-0100
670557-0100
680557-0100
690557-0100
700557-0100
710557-0100
720557-0100
730557-0100
740557-0100
750557-0100
760557-0100
770557-0100
780557-0100
790557-0100
800557-0100
810557-0100
820557-0100
830557-0100
840557-0100
850557-0100
860557-0100
870557-0100
880557-0100
890557-0100
900557-0100
910557-0100
920557-0100
930557-0100
940557-0100
950557-0100
960557-0100
970557-0100
980557-0100
990557-0100
100557-0100
101557-0100
102557-0100
103557-0100
104557-0100
105557-0100
106557-0100
107557-0100
108557-0100
109557-0100
110557-0100
111557-0100
112557-0100
113557-0100
114557-0100
115557-0100
116557-0100
117557-0100
118557-0100
119557-0100
120557-0100
121557-0100
122557-0100
123557-0100
124557-0100
125557-0100
126557-0100
127557-0100
128557-0100
129557-0100
130557-0100
131557-0100
132557-0100
133557-0100
134557-0100
135557-0100
136557-0100
137557-0100
138557-0100
139557-0100
140557-0100
141557-0100
142557-0100
143557-0100
144557-0100
145557-0100
146557-0100
147557-0100
148557-0100
149557-0100
150557-0100
151557-0100
152557-0100
153557-0100
154557-0100
155557-0100
156557-0100
157557-0100
158557-0100
159557-0100
160557-0100
161557-0100
162557-0100
163557-0100
164557-0100
165557-0100
166557-0100
167557-0100
168557-0100
169557-0100
170557-0100
171557-0100
172557-0100
173557-0100
174557-0100
175557-0100
176557-0100
177557-0100
178557-0100
179557-0100
180557-0100
181557-0100
182557-0100
183557-0100
184557-0100
185557-0100
186557-0100
187557-0100
188557-0100
189557-0100
190557-0100
191557-0100
192557-0100
193557-0100
194557-0100
195557-0100
196557-0100
197557-0100
198557-0100
199557-0100
200557-0100
201557-0100
202557-0100
203557-0100
204557-0100
205557-0100
206557-0100
207557-0100
208557-0100
209557-0100
210557-0100
211557-0100
212557-0100
213557-0100
214557-0100
215557-0100
216557-0100
217557-0100
218557-0100
219557-0100
220557-0100
221557-0100
222557-0100
223557-0100
224557-0100
225557-0100
226557-0100
227557-0100
228557-0100
229557-0100
230557-0100
231557-0100
232557-0100
233557-0100
234557-0100
235557-0100
236557-0100
237557-0100
238557-0100
239557-0100
240557-0100
241557-0100
242557-0100
243557-0100
244557-0100
245557-0100
246557-0100
247557-0100
248557-0100
249557-0100
250557-0100
251557-0100
252557-0100
253557-0100
254557-0100
255557-0100
256557-0100
257557-0100
258557-0100
259557-0100
260557-0100
261557-0100
262557-0100
263557-0100
264557-0100
265557-0100
266557-0100
267557-0100
268557-0100
269557-0100
270557-0100
271557-0100
272557-0100
273557-0100
274557-0100
275557-0100
276557-0100
277557-0100
278557-0100
279557-0100
280557-0100
281557-0100
282557-0100
283557-0100
284557-0100
285557-0100
286557-0100
287557-0100
288557-0100
289557-0100
290557-0100
291557-0100
292557-0100
293557-0100
294557-0100
295557-0100
296557-0100
297557-0100
298557-0100
299557-0100
300557-0100
301557-0100
302557-0100
303557-0100
304557-0100
305557-0100
306557-0100
307557-0100
308557-0100
309557-0100
310557-0100
311557-0100
312557-0100
313557-0100
314557-0100
315557-0100
316557-0100
317557-0100
318557-0100
319557-0100
320557-0100
321557-0100
322557-0100
323557-0100
324557-0100
325557-0100
326557-0100
327557-0100
328557-0100
329557-0100
330557-0100
331557-0100
332557-0100
333557-0100
334557-0100
335557-0100
336557-0100
337557-0100
338557-0100
339557-0100
340557-0100
341557-0100
342557-0100
343557-0100
344557-0100
345557-0100
346557-0100
347557-0100
348557-0100
349557-0100
350557-0100
351557-0100
352557-0100
353557-0100
354557-0100
355557-0100
356557-0100
357557-0100
358557-0100
359557-0100
360557-0100
361557-0100
362557-0100
363557-0100
364557-0100
365557-0100
366557-0100
367557-0100
368557-0100
369557-0100
370557-0100
371557-0100
372557-0100
373557-0100
374557-0100
375557-0100
376557-0100
377557-0100
378557-0100
379557-0100
380557-0100
381557-0100
382557-0100
383557-0100
384557-0100
385557-0100
386557-0100
387557-0100
388557-0100
389557-0100
390557-0100
391557-0100
392557-0100
393557-0100
394557-0100
395557-0100
396557-0100
397557-0100
398557-0100
399557-0100
400557-0100
401557-0100
402557-0100
403557-0100
404557-0100
405557-0100
406557-0100
407557-0100
408557-0100
409557-0100
410557-0100
411557-0100
412557-0100
413557-0100
414557-0100
415557-0100
416557-0100
417557-0100
418557-0100
419557-0100
420557-0100
421557-0100
422557-0100
423557-0100
424557-0100
425557-0100
426557-0100
427557-0100
428557-0100
429557-0100
430557-0100
431557-0100
432557-0100
433557-0100
434557-0100
435557-0100
436557-0100
437557-0100
438557-0100
439557-0100
440557-0100
441557-0100
442557-0100
443557-0100
444557-0100
445557-0100
446557-0100
447557-0100
448557-0100
449557-0100
450557-0100
451557-0100
452557-0100
453557-0100
454557-0100
455557-0100
456557-0100
457557-0100
458557-0100
459557-0100
460557-0100
461557-0100
462557-0100
463557-0100
464557-0100
465557-0100
466557-0100
467557-0100
468557-0100
469557-0100
470557-0100
471557-0100
472557-0100
473557-0100
474557-0100
475557-0100
476557-0100
477557-0100
478557-0100
479557-0100
480557-0100
481557-0100
482557-0100
483557-0100
484557-0100
485557-0100
486557-0100
487557-0100
488557-0100
489557-0100
490557-0100
491557-0100
492557-0100
493557-0100
494557-0100
495557-0100
496557-0100
497557-0100
498557-0100
499557-0100
500557-0100
501557-0100
502557-0100
503557-0100
504557-0100
505557-0100
506557-0100
507557-0100
508557-0100
509557-0100
510557-0100
511557-0100
512557-0100
513557-0100
514557-0100
515557-0100
516557-0100
517557-0100
518557-0100
519557-0100
520557-0100
521557-0100
522557-0100
523557-0100
524557-0100
525557-0100
526557-0100
527557-0100
528557-0100
529557-0100
530557-0100
531557-0100
532557-0100
533557-0100
534557-0100
535557-0100
536557-0100
537557-0100
538557-0100
539557-0100
540557-0100
541557-0100
542557-0100
543557-0100
544557-0100
545557-0100
546557-0100
547557-0100
548557-0100
549557-0100
550557-0100
551557-0100
552557-0100
553557-0100
554557-0100
555557-0100
556557-0100
557557-0100
558557-0100
559557-0100
560557-0100
561557-0100
562557-0100
563557-0100
564557-0100
565557-0100
566557-0100
567557-0100
568557-0100
569557-0100
570557-0100
571557-0100
572557-0100
573557-0100
574557-0100
575557-0100
576557-0100
577557-0100
578557-0100
579557-0100
580557-0100
581557-0100
582557-0100
583557-0100
584557-0100
585557-0100
586557-0100
587557-0100
588557-0100
589557-0100
590557-0100
591557-0100
592557-0100
593557-0100
594557-0100
595557-0100
596557-0100
597557-0100
598557-0100
599557-0100
600557-0100
601557-0100
602557-0100
603557-0100
604557-0100
605557-0100
606557-0100
607557-0100
608557-0100
609557-0100
610557-0100
611557-0100
612557-0100
613557-0100
614557-0100
615557-0100
616557-0100
617557-0100
618557-0100
619557-0100
620557-0100
621557-0100
622557-0100
623557-0100
624557-0100
625557-0100
626557-0100
627557-0100
628557-0100
629557-0100
630557-0100
631557-0100
632557-0100
633557-0100
634557-0100
635557-0100
636557-0100
637557-0100
638557-0100
639557-0100
640557-0100
641557-0100
642557-0100
643557-0100
644557-0100
645557-0100
646557-0100
647557-0100
648557-0100
649557-0100
650557-0100
651557-0100
652557-0100
653557-0100
654557-0100
655557-0100
656557-0100
657557-0100
658557-0100
659557-0100
660557-0100
661557-0100
662557-0100
663557-0100
664557-0100
665557-0100
666557-0100
667557-0100
668557-0100
669557-0100
670557-0100
671557-0100
672557-0100
673557-0100
674557-0100
675557-0100
676557-0100
677557-0100
678557-0100
679557-0100
680557-0100
681557-0100
682557-0100
683557-0100
684557-0100
685557-0100
686557-0100
687557-0100
688557-0100
689557-0100
690557-0100
691557-0100
692557-0100
693557-0100
694557-0100
695557-0100
696557-0100
697557-0100
698557-0100
699557-0100
700557-0100
701557-0100
702557-0100
703557-0100
704557-0100
705557-0100
706557-0100
707557-0100
708557-0100
709557-0100
710557-0100
711557-0100
712557-0100
713557-0100
714557-0100
715557-0100
716557-0100
717557-0100
718557-0100
719557-0100
720557-0100
721557-0100
722557-0100
723557-0100
724557-0100
725557-0100
726557-0100
727557-0100
728557-0100
729557-0100
730557-0100
731557-0100
732557-0100
733557-0100
734557-0100
735557-0100
736557-0100
737557-0100
738557-0100
739557-0100
740557-0100
741557-0100
742557-0100
743557-0100
744557-0100
745557-0100
746557-0100
747557-0100
748557-0100
749557-0100
750557-0100
751557-0100
752557-0100
753557-0100
754557-0100
755557-0100
756557-0100
757557-0100
758557-0100
759557-0100
760557-0100
761557-0100
762557-0100
763557-0100
764557-0100
765557-0100
766557-0100
767557-0100
768557-0100
769557-0100
770557-0100
771557-0100
772557-0100
773557-0100
774557-0100
775557-0100
776557-0100
777557-0100
778557-0100
779557-0100
780557-0100
781557-0100
782557-0100
783557-0100
784557-0100
785557-0100
786557-0100
787557-0100
788557-0100
789557-0100
790557-0100
791557-0100
792557-0100
793557-0100
794557-0100
795557-0100
796557-0100
797557-0100
798557-0100
799557-0100
800557-0100
801557-0100
802557-0100
803557-0100
804557-0100
805557-0100
806557-0100
807557-0100
808557-0100
809557-0100
810557-0100
811557-0100
812557-0100
813557-0100
814557-0100
815557-0100
816557-0100
817557-0100
818557-0100
819557-0100
820557-0100
821557-0100
822557-0100
823557-0100
824557-0100
825557-0100
826557-0100
827557-0100
828557-0100
829557-0100
830557-0100
831557-0100
832557-0100
833557-0100
834557-0100
835557-0100
836557-0100
837557-0100
838557-0100
839557-0100
840557-0100
841557-0100
842557-0100
843557-0100
844557-0100
845557-0100
846557-0100
847557-0100
848557-0100
849557-0100
850557-0100
851557-0100
852557-0100
853557-0100
854557-0100
855557-0100
856557-0100
857557-0100
858557-0100
859557-0100
860557-0100
861557-0100
862557-0100
863557-0100
864557-0100
865557-0100
866557-0100
867557-0100
868557-0100
869557-0100
870557-0100
871557-0100
872557-0100
873557-0100
874557-0100
875557-0100
876557-0100
877557-0100
878557-0100
879557-0100
880557-0100
881557-0100
882557-0100
883557-0100
884557-0100
885557-0100
886557-0100
887557-0100
888557-0100
889557-0100
890557-0100
89155

cooled mixture is added an appropriate agent for effecting the primer extension reaction (called herein "agent for polymerization"), and the reaction is allowed to occur under conditions known in the art. The agent for polymerization may also be added together with the other reagents if it is heat stable. This synthesis (or amplification) reaction may occur at room temperature up to a temperature above which the agent for polymerization no longer functions. Thus, for example, if DNA polymerase is used as the agent, the temperature is generally no greater than about 40°C. Most conveniently the reaction occurs at room temperature.

[0072] The agent for polymerization may be any compound or system that will function to accomplish the synthesis of primer extension products, including enzymes. Suitable enzymes for this purpose include, for example, E. coli DNA polymerase I, Klenow fragment of E. coli DNA polymerase I, T4 DNA polymerase, other available DNA polymerases, polymerase muteins, reverse transcriptase, and other enzymes, including heat-stable enzymes (i.e., those enzymes which perform primer extension after being subjected to temperatures sufficiently elevated to cause denaturation such as Taq DNA polymerase, and the like). Suitable enzymes will facilitate combination of the nucleotides in the proper manner to form the primer extension products that are complementary to each locus nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths. There may be agents for polymerization, however, which initiate synthesis at the 5' end and proceed in the other direction, using the same process as described above.

[0073] Preferably, the method of amplifying is by PCR, as described herein and as is commonly used by those of ordinary skill in the art. However, alternative methods of amplification have been described and can also be employed. PCR techniques and many variations of PCR are known. Basic PCR techniques are described by Saiki *et al.* (1988 Science 239:487-491) and by U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference.

[0074] The conditions generally required for PCR include temperature, salt, cation, pH and related conditions needed for efficient copying of the master-cut fragment. PCR

1005979-0400

conditions include repeated cycles of heat denaturation (i.e. heating to at least about 95° C.) and incubation at a temperature permitting primer: adaptor hybridization and copying of the master-cut DNA fragment by the amplification enzyme. Heat stable amplification enzymes like the pwo, *Thermus aquaticus* or *Thermococcus litoralis* DNA polymerases which eliminate the need to add enzyme after each denaturation cycle, are commercially available. The salt, cation, pH and related factors needed for enzymatic amplification activity are available from commercial manufacturers of amplification enzymes.

[0075] As provided herein an amplification enzyme is any enzyme which can be used for in vitro nucleic acid amplification, e.g. by the above-described procedures. Such amplification enzymes include pwo, *Escherichia coli* DNA polymerase I, Klenow fragment of *E. coli* DNA polymerase I, T4 DNA polymerase, T7 DNA polymerase, *Thermus aquaticus* (Taq) DNA polymerase, *Thermococcus litoralis* DNA polymerase, SP6 RNA polymerase, T7 RNA polymerase, T3 RNA polymerase, T4 polynucleotide kinase, Avian Myeloblastosis Virus reverse transcriptase, Moloney Murine Leukemia Virus reverse transcriptase, T4 DNA ligase, *E. coli* DNA ligase or Q β replicase. Preferred amplification enzymes are the pwo and Taq polymerases. The pwo enzyme is especially preferred because of its fidelity in replicating DNA.

[0076] Once amplified, the nucleic acid can be attached to a solid support, such as a membrane, and can be hybridized with any probe of interest, to detect any nucleic acid sequence. Several membranes are known to one of skill in the art for the adhesion of nucleic acid sequences. Specific non-limiting examples of these membranes include nitrocellulose (NITROPURE®) or other membranes used in for detection of gene expression such as polyvinylchloride, diazotized paper and other commercially available membranes such as GENESCREEN®, ZETAPROBE® (Biorad), and NYTRAN®. Methods for attaching nucleic acids to these membranes are well known to one of skill in the art. Alternatively, screening can be done in a liquid phase.

[0077] In nucleic acid hybridization reactions, the conditions used to achieve a particular level of stringency will vary, depending on the nature of the nucleic acids being hybridized. For example, the length, degree of complementarity, nucleotide sequence composition (e.g.,

10059573-01200

GC v. AT content), and nucleic acid type (e.g., RNA v. DNA) of the hybridizing regions of the nucleic acids can be considered in selecting hybridization conditions. An additional consideration is whether one of the nucleic acids is immobilized, for example, on a filter.

[0078] An example of progressively higher stringency conditions is as follows: 2 x SSC/0.1% SDS at about room temperature (hybridization conditions); 0.2 x SSC/0.1% SDS at about room temperature (low stringency conditions); 0.2 x SSC/0.1% SDS at about 42°C (moderate stringency conditions); and 0.1 x SSC at about 68°C (high stringency conditions). Washing can be carried out using only one of these conditions, e.g., high stringency conditions, or each of the conditions can be used, e.g., for 10-15 minutes each, in the order listed above, repeating any or all of the steps listed. However, as mentioned above, optimal conditions will vary, depending on the particular hybridization reaction involved, and can be determined empirically. In general, conditions of high stringency are used for the hybridization of the probe of interest.

[0079] The probe of interest can be detectably labeled, for example, with a radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound, a metal chelator, or an enzyme. Those of ordinary skill in the art will know of other suitable labels for binding to the probe, or will be able to ascertain such, using routine experimentation.

[0080] Another embodiment of the invention provides a method of determining a predisposition to a cellular proliferative disorder of breast tissue in a subject comprising determining the state of methylation of one or more nucleic acids isolated from the subject, wherein the nucleic acid is selected from the group consisting of Twist, cyclin D2, RAR β 2, HOXA5, WT1, 14.3.3 sigma, estrogen receptor, NES-1, RASSF1A, HIN-1, and combinations thereof; and wherein the state of methylation of one or more nucleic acids as compared with the state of methylation of said nucleic acid from a subject not having a predisposition to the cellular proliferative disorder of breast tissue is indicative of a cell proliferative disorder of breast tissue in the subject.

4005579-013803

[0081] As used herein, "predisposition" refers to an increased likely that an individual will have a disorder. Although a subject with a predisposition does not yet have the disorder, there exists an increased propensity to the disease.

[0082] Another embodiment of the invention provides a method for diagnosing a cellular proliferative disorder of breast tissue in a subject comprising contacting a nucleic acid-containing specimen from the subject with an agent that provides a determination of the methylation state of nucleic acids in the specimen, and identifying the methylation state of at least one region of least one nucleic acid, wherein the methylation state of at least one region of at least one nucleic acid that is different from the methylation state of the same region of the same nucleic acid in a subject not having the cellular proliferative disorder is indicative of a cellular proliferative disorder of breast tissue in the subject.

[0083] Invention methods are ideally suited for the preparation of a kit. Therefore, in accordance with another embodiment of the present invention, there is provided a kit it useful for the detection of a cellular proliferative disorder in a subject. Invention kits include a carrier means compartmentalized to receive a sample therein, one or more containers comprising a first container containing a reagent which modifies unmethylated cytosine and a second container containing primers for amplification of a CpG-containing nucleic acid, wherein the primers distinguish between modified methylated and nonmethylated nucleic acid. Primers contemplated for use in accordance with the invention include those set forth in SEQ ID NOs: 7-14, 21-24, 37-40, 49-64, 69-72, 77-80, 85-90, 116-119, 124-128, and combinations thereof.

[0084] Carrier means are suited for containing one or more container means such as vials, tubes, and the like, each of the container means comprising one of the separate elements to be used in the method. In view of the description provided herein of invention methods, those of skill in the art can readily determine the apportionment of the necessary reagents among the container means. For example, one of the container means can comprise a container containing an oligonucleotide for ligation to nucleic acid cleaved by a methylation sensitive restriction endonuclease. One or more container means can also be included comprising a primer complementary to the oligonucleotide. In addition, one or more container means can

10050579-010001

also be included which comprise a methylation sensitive restriction endonuclease. One or more container means can also be included containing an isoschizomer of said methylation sensitive restriction enzyme.

[0085] The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples, which are provided herein for purposes of illustration only and are not intended to limit the scope of the invention.

[0086] Although the invention has been described with reference to the presently preferred embodiment, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

EXAMPLE 1

Methylation Status of Wilms' tumor suppressor gene (WT1)

[0087] The extent of methylation of the WT1-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors, and expression of the WT1 mRNA and protein in the same cells and tissues was examined.

[0088] **Cell lines and finite life span cultures** Cell lines were obtained from ATCC (Rockville, MD) and grown according to conditions specified. Also utilized were three independent cultures of finite life span human mammary epithelial cells (HMEC): 16637 (Clonetics, Walkersville, MD) and 1-26, 3-14 (kindly provided by Dr. Steve Ethier, Univ. Michigan, Ann Arbor, MI). When indicated, cell lines were treated with 0.75 μ M 5-aza-2'-deoxycytidine (5-aza-dC) or with 100 ng/ml Trichostatin A (TSA) as described in Ferguson, *et al.* (*Proc Natl Acad Sci U S A.* (2000) 97:6049-54).

[0089] **Tumors and Organoids** Primary breast tumors were obtained from the Johns Hopkins frozen tumor bank. Mammary organoids were prepared from reduction mammoplasty specimens of women with benign (B) or no (N) abnormalities in the breast as

2008270-6755007 10055579-012802

described in Fujii, *et al.* (Oncogene. 16: 2159-64, 1998). Briefly, the specimens were enzymatically digested into duct-like structures (organoids), filtered, histologically confirmed to contain more than 80% epithelial cells, and frozen at -70° C until used. Also utilized were highly purified myo- and luminal epithelial cells isolated by differential centrifugation and fluorescence-activated cell sorting of enzymatically digested normal mammary specimens (Gomm, *et al.*, (1995) *Anal Biochem.* 226 91-9).

[0090] RT-PCR for WT1 mRNA Methods for RNA extraction and RT-PCR are known to those of skill in the art. The sequences of the primers used are as follows: for amplifying the 555 bp region surrounding WT1 exon 5, 5'-GCGGCGCAGTTCCCAACCA-3' (sense, nucleotides 882-901; SEQ ID NO:1) and 5'-ATGTTTCTCACCAGTGTGCTT-3' (antisense, nucleotides 1416-1437; SEQ ID NO:2); for amplifying the 382 bp region surrounding the KTS insert, 5'-GCATCTGAAACCAGTGAGAA-3' (sense, nucleotides 1320-1339; SEQ ID NO:3) and 5'-TTTCTCTGATGCATGTTG-3' (antisense, nucleotides 1685-1702; SEQ ID NO:4). Amplification was performed using a hot-start protocol: samples were heated to 94°C for 4 minutes and then cooled to 80°C prior to the addition of Taq polymerase (RedTaq, Sigma, St. Louis, MO). Samples were then heated to 94°C for 30 seconds followed by either 50°C for 30 seconds (for the KTS primers) or 56°C for 30 seconds (for the Exon 5 primers) and then 72°C for 1 minute for 40 cycles. PCR products were resolved by electrophoresis, using a 2% agarose gel for the exon 5-splice variants and a 12% polyacrylamide gel to resolve the KTS insert variants. Co-amplification of the ribosomal RNA 36B4 was performed as an internal control using the following primers: 5'-GATTGGCTACCCAACTGTTGCA-3' (sense; SEQ ID NO:5) and 5'-CAGGGGCAGCAGCCACAAAGGC-3' (antisense; SEQ ID NO:6).

[0091] Northern blots Total RNA was extracted as described above. After electrophoresis through a 1.5% agarose gel in MOPS buffer with 6.7% formaldehyde, RNA was transferred to nitrocellulose. Blots were probed with a PCR product corresponding to the WT1 zinc finger region, amplified using the primers described above and labeled by random priming using standard techniques.

105579-01200

[0092] Methylation-specific PCR Genomic DNA was isolated using standard techniques and treated with sodium bisulfite as described elsewhere (Herman, *et al.*, Proc Natl Acad Sci U S A. 93: 9821-6, 1996). Methylation-specific PCR was performed using the following primers: to detect methylated promoter DNA, 5'-TTTGGGTAAAGTTAGGCGTCGTCG-3' (sense; SEQ ID NO:7) and 5'-ACACTACTCCTCGTACGACTCCG-3' (antisense; SEQ ID NO:8); to detect unmethylated promoter DNA, 5'-TTTGGGTAAAGTTAGGTGTTGTTG-3' (sense; SEQ ID NO:9) and 5'-ACACTACTCCTCATACAACTCCA-3' (antisense; SEQ ID NO:10); to detect methylated intron 1 DNA, 5'-CGTCGGGTGAAGGCGGGTAAT-3' (sense; SEQ ID NO:11) and 5'-CGAACCCGAACCTACGAAACC-3' (antisense; SEQ ID NO:12); to detect unmethylated intron 1 DNA, 5'-TGTTGGGTGAAGGTGGGTAAT-3' (sense; SEQ ID NO:13) and 5'-CAAACCCAAACCTACAAAACC-3' (antisense; SEQ ID NO:14). The PCR reaction was as described above, except that the annealing temperature was 59°C, and the extension time was 45 seconds.

[0093] Western blots Total protein from cell lines was obtained from material harvested in TriReagent (Molecular Research Center, Cincinnati, OH) and initially used for RNA isolation. Protein purification was according to the manufacturer's protocol. After separation by SDS-PAGE and electrophoretic transfer to nitrocellulose membranes, proteins were incubated with an anti-WT1 antibody [WT (C-19); sc-192, Santa Cruz Biotechnology, Santa Cruz, CA] diluted 1:1000 in the blocking solution. Horseradish peroxidase-conjugated antibody against rabbit IgG (Amersham, Arlington Heights, IL) was used at 1:1000, and binding was revealed using enhanced chemiluminescence (Amersham, Arlington Heights, IL).

[0094] Expression of WT1 mRNA in mammary epithelial and breast cancer cell lines To evaluate WT1 expression in the breast, mRNA expression was analyzed by RT-PCR in a panel of normal and transformed cell lines. No WT1 RNA was detected in 3 independently derived finite lifespan mammary epithelial strains: HMEC 16637, 1-26, and 3-14. Among the three immortal breast epithelial cell lines, WT1 expression was observed in HMECs HBL-100 and MCF-10A, but not in H16N. WT1 mRNA expression was examined in nine breast cancer cell lines, expression was easily detectable in five: HS578T, T47D,

10059579-010003

MDA-MB-468, 21MT, and 21PT, and undetectable in the remaining four: SKBR3, MDA-MB-435, MCF-7 and MDA-MB-231.

[0095] The specific expression of WT1 isoforms lacking the fifth exon and lacking the KTS insert has been reported to occur in breast cancer (Silberstein, *et al.*, Proc Natl Acad Sci U S A. 94: 8132-7, 1997). To determine if differential expression of WT1 splice variants is seen in breast cancer cell lines, PCR primers were designed spanning the fifth exon such that mRNA encoding the isoform containing exon 5 yielded a 555 bp PCR product, while if exon 5 were missing a 504 bp PCR product was generated. PCR primers spanning the region of the KTS insert, such that an mRNA containing the insert would yield a 382 bp product, while an RNA lacking the insert would generate a 373 bp product were also used. Contrary to the findings in the published report (Silberstein *et al.*, *supra*), in the five WT1-expressing breast cancer cell lines, and in the WT1-expressing immortalized HMECs, all four splice variants- the two exon 5 isoforms and the two KTS isoforms, were present.

[0096] To confirm these results, Northern blot analysis was performed using total RNA isolated from a number of breast cancer cell lines. Similar to the results obtained by RT-PCR, WT1 mRNA expression was readily detected in HBL-100, HS-578T, T47D, and MDA-MB-468 cells but was not detected in MDA-MB-435, MDA-MB-231, SKBR3, or MCF-7 cells.

[0097] Thus, WT1 mRNA expression was undetectable in finite life span primary breast epithelial cell cultures, but was easily detectable in the neoplastic and immortalized HMECs and in seven of twelve breast cancer cell lines. Also, the striking correlation between results from Northern blots and RT-PCR experiments validated the RT-PCR protocol for the detection of WT1 mRNA expression.

[0098] **Methylation of the WT1 locus in breast cancer cell lines** The promoter and first intron of the WT1 gene contain dense CpG islands. These sequence elements are frequently sites of DNA methylation, and play a role in transcriptional silencing (Nan, *et al.*, Cell. 88: 471-81, 1997; Ng, *et al.*, Nat Genet. 23: 58-61, 1999). To determine whether methylation silences gene expression in the WT1-negative cell lines, the status of the WT1 promoter in

1055579.010002

the breast cancer cell lines was investigated. The promoter was methylated in the 4 cell lines that did not express WT1, but not in the 5 cell lines that did, consistent with the idea that methylation is a critical determinant of WT1 expression. There was one exception to this correlation. T47D cells contained methylated WT1 sequences but nevertheless expressed WT1 mRNA, suggesting that, in this case, methylation alone is insufficient to silence expression.

[0099] Promoter methylation is postulated to silence transcription, at least in part, by recruitment of histone deacetylase (HDAC) to hypermethylated loci (Nan, *et al. supra* and Ng, *et al., supra*). In order to assess the functional significance of WT1 promoter methylation, MDA-MB-231 and MCF-7 cells were treated with 5-aza-deoxyC, an inhibitor of DNA methyltransferases, or with TSA, an inhibitor of HDAC. As demonstrated before (Laux, *et al.*, Breast Cancer Res Treat. 56: 35-43, 1999), treatment with 5-aza-deoxyC resulted in WT1 expression by MDA-MB-231 cells. Interestingly, this treatment did not cause WT1 expression in MCF-7 cells, nor did TSA restore expression in either cell line. In the same samples, these treatments restored expression of 14.3.3 σ (Ferguson, *et al.*, *Proc Natl Acad Sci U S A.* (2000) 97:6049-54). These findings suggest that while promoter methylation correlates with gene silencing, it may not play a causal role.

[0100] **Expression of WT1 in primary breast tissue** These findings from cell lines were expanded to patient samples, including normal breast epithelium and primary breast tumors. Breast carcinomas arise from luminal epithelial cells in the mammary duct. Normal breast tissue also contains a layer of myoepithelial cells, which overlie the luminal epithelium. To ensure that the normal samples contained luminal epithelial cells, three different types of epithelial cell preparations were used, including (1) three short term cultures of HMECs, (2) nine organoid preparations of mammary ducts, and (3) eight samples of highly purified luminal and myoepithelial cells (isolated from 4 patient samples).

[0101] WT1 expression was not detected by RT-PCR in 3 HMEC samples, in eight out of nine breast organoid preparations, nor in any of eight purified epithelial cell preparations. By western blotting, WT1 protein was not detected in three organoid samples nor in two

10055579.042007

HMECs. In contrast, WT1 expression was easily detectable in 27 out of 31 (87%) primary breast carcinomas.

[0102] The HMECs did not express WT1; however, RT-PCR using primers described above demonstrated the expression of Exon 5 (+) and Exon 5 (-) isoforms in five out of seven tumors, while the remaining two expressed only the Exon 5 (+) isoform. KTS (+) and KTS (-) isoforms were detected in all nine tumors examined. Thus, a majority of the tumors expressed both Exon 5 splice variants of WT1, and all of the tumors express both splice variants involving the KTS insert. Interestingly, the sole breast organoid sample that expressed WT1 expressed all four splice variants as well.

[0103] **Methylation of WT1-associated CpG islands in normal and malignant breast tissue** Since methylation of the promoter-associated CpG island correlated with a lack of WT1 expression in breast cancer cell lines, the methylation status of the promoter and first intron CpG islands was studied in this panel of breast organoids and carcinomas. Prior studies demonstrating tumor-specific methylation of the CpG islands associated with the WT1 gene have employed methylation-sensitive restriction enzymes (Huang, *et al.*, Cancer Res. 57: 1030-4, 1997; Laux, *et al.*, Breast Cancer Res Treat. 56: 35-43, 1999; and Huang, *et al.*, Hum Mol Genet. 8: 459-70, 1999). This technique is a reliable way to identify individual methylated sites, but it is unable to assess large-scale methylation patterns. The density of methylation, rather than methylation of any specific CpG dinucleotide, is responsible for gene silencing (Herman, *et al.*, Semin Cancer Biol. 9: 359-67, 1999). Therefore, methylation of the CpG islands was evaluated using methylation specific PCR (MSP). This method allows the direct evaluation of several methylation sites per PCR reaction, and choosing a variety of sequences for PCR primers allows the rapid assessment of many CpG dinucleotides (Herman, *et al.*, Proc Natl Acad Sci U S A. 93: 9821-6, 1996).

[0104] MSP was performed using DNA extracted from 19 primary tumors and nine breast organoid preparations. The WT1 promoter CpG island was unmethylated in DNA from all nine organoid samples. In contrast, six of 19 tumors contained methylated DNA, and the remaining 13 were completely unmethylated. This rate of promoter methylation (32%) is not dissimilar to the 25% incidence reported by Laux *et al.* (Breast Cancer Res Treat. 56: 35-43,

105579-3100

1999). Thus, methylation of the WT1 promoter is a tumor-specific phenomenon. Contrary to expectation, however, each of the six tumors that contained methylated WT1 also expressed WT1 protein. WT1 gene methylation, therefore, was not effective in silencing gene expression. Next, the CpG island in the first intron of the WT1 gene, a region where tumor-specific methylation has also been previously reported was examined. Methylation of WT1 was detected in all three breast organoid preparations and in nine of ten tumor samples evaluated. Thus, the first intron of WT1 is methylated in both normal and malignant breast tissue, and is unrelated to tumorigenesis.

[0105] Methylation of the CpG island associated with the WT1 promoter is associated with a gene silencing in several breast cancer cell lines. While treatment of MDA-MB-231 cells with the methyltransferase inhibitor 5-aza-deoxycytidine results in re-expression of the gene, this was not seen in MCF-7 cells. Additionally, treatment with the HDAC inhibitor TSA had no effect on WT1 expression, suggesting that DNA methylation and histone acetylation play only minor roles in the regulation of WT1 expression in mammary epithelium.

[0106] This study demonstrates tumor-specific methylation of the CpG islands of WT1. Surprisingly, expression of WT1 mRNA and protein in the majority of breast cancer samples evaluated was also found, including in every sample that contained methylated DNA. These findings that breast carcinomas express WT1 despite tumor-specific gene methylation emphasizes the importance of evaluating methylation and gene expression concurrently in the same tissue.

[0107] WT1 mRNA was readily detected in tumor samples using a single step PCR protocol. While it is possible to detect WT1 expression in normal epithelium using a nested PCR, this would not alter the finding that the gene is overexpressed in tumors compared with normal tissue. The use of RT-PCR may allow the detection of a relatively weakly expressed gene, but WT1 protein was readily detected by Western blotting in tumors. Since protein is the functional species, this finding suggests that WT1 is abundant enough in tumors to play a functional role.

10056575:042808

[0108] These data also reveal a discrepancy between gene regulation in tissue culture and *in vivo*. Methylation of the WT1 promoter is associated with gene silencing in breast cancer cell lines. In contrast, the promoter-associated CpG island was methylated in 32% of the tumors examined; contrary to expectation, these tumors express WT1. These data highlight the fact that there are multiple mechanisms for gene silencing, of which hypermethylation of a CpG island is only one. More importantly, these findings emphasize the idea that cell lines do not necessarily reflect the *in vivo* situation. They also serve to point out that hypermethylation of a CpG island may be insufficient to silence expression, demonstrating the importance of assessing gene expression as well as promoter methylation status when evaluating the role of a particular gene in a particular tumor type.

[0109] In summary, these data demonstrate that WT1 is not expressed in normal breast epithelium and is over-expressed in the majority of primary breast tumors. Tumor-specific methylation of the CpG island occurs in breast cancer, but appears to be inconsequential to gene expression.

EXAMPLE 2

Hypermethylation and loss of expression of cyclin D2

[0110] The extent of methylation of the cyclin D2-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors, and expression of the cyclin D2 mRNA and protein in the same cells and tissues was examined.

[0111] **Cell Lines and Tissues** The breast cancer cell lines MDAMB435, MCF7, T47D, SKBR3, ZR75.1, MDAMB468, HS578T, MDAMB231 and the immortal human mammary epithelial cell lines (HMEC) MCF10A and HBL100 were obtained and maintained in culture according to instructions (ATCC, Rockville, MD). The two matched tumor cell lines, 21PT, derived from a primary tumor and 21MT, from the metastasis of the same patient, were propagated as described elsewhere. The breast cancer cell line, MW, was obtained from Dr. Renato Dulbecco. HMEC-H16N (immortalized with HPV) was kindly provided by Dr. Vimla Band. Cultured finite life span human breast epithelial cell strains 04372, 219-6,

and 166372 were obtained from Clonetics (Walkersville, MD), and HMEC strains 1-26 and 3-14 were kindly provided by Dr. Steve Ethier. Finite life span HMEC 184, the immortalized HMECs 184A1 (passage 15 and 99) and 184B5 were kindly provided by Dr. Martha Stampfer, and grown as described on the worldwide web site lbl.gov/LBL-Programs/mrgs/review.html. Cell extracts from finite lifespan HMECs 70N and 81N were kindly provided by Dr. Khandan Keyomarsi. Mammary organoids were prepared from reduction mammoplasty specimens of women with benign or no abnormalities in the breast following collagenase digestion as described in Bergstraesser LM, (1993). Human mammary luminal and myoepithelial cells were prepared by progressive collagenase digestion of breast tissue, sedimentated to obtain organoids (ductal and lobulo-alveolar fragments), cultured short term, and finally highly enriched by using an immunomagnetic separation technique (Niranjan B, 1995).

[0112] Primary breast tumor tissues were obtained after surgical resection at the Johns Hopkins University and Duke University, and stored frozen at -80°C . Samples containing greater than 50% tumor cells were selected following microscopic examination of representative tissue sections from each tumor. Microdissection of carcinoma and ductal carcinomas *in situ* (DCIS) lesions from eight micron cryosections was performed by using a laser capture microscope, or by manually scraping the cells with a 25G needle under 40X magnification. Genomic DNA was extracted by incubating the microdissected cells at 55°C x12 h in 50 μl buffer containing 10 mM Tris Cl (pH 8.0), 1 mM EDTA, 0.1% Tween 20, and 0.5 $\mu\text{g}/\mu\text{l}$ proteinase K. The extract was heat inactivated at 95°C for 5 min., and used directly for sodium bisulfite treatment.

[0113] RT-PCR RNA was treated with RNase-free DNase (Boehringer-Mannheim) (0.5-1u/ μl) for 30 min. at 37°C , followed by heat inactivation at 65°C for 10 min. RT reactions contained 2 μg DNase treated RNA, 0.25 $\mu\text{g}/\mu\text{l}$ pdN6 random primers (Pharmacia), 1X first strand buffer (GibcoBRL), 1 mM dNTP (Pharmacia), and 200 U MMLV-RT (GibcoBRL), and were incubated for 1h at 37°C followed by heat inactivation at 75°C for 5 min. PCR was performed using the primers 5'-CATGGAGCTGCTGTGCCACG-3' (sense; SEQ ID NO:15) and 5'-CCGACCTACCTCCAGCATCC-3' (antisense; SEQ ID NO:16) for

SECRET

SECRET

SECRET

SECRET

NaCl and PMSF, and sonicated. Twenty µg of protein were fractionated on 12.5% SDS-PAGE and transferred by electrophoresis to a nylon membrane. The blot was incubated with anti-cyclin D2 antibody (Ab-4, "cocktail" mouse monoclonal antibodies, Neomarkers, San Diego, CA) diluted 1:200 in 5% skim milk, for 2h at room temperature. Horseradish peroxidase-conjugated antibody anti-mouse IgG (Amersham) was used at 1:1000, and binding was revealed using enhanced chemiluminescence (Amersham).

[0117] Cyclin D2 mRNA expression in breast cancer Serial analysis of gene expression (SAGE) and subsequent microarray analysis previously revealed that, compared with finite lifespan HMECs, cyclin D2 expression was significantly lower in a small panel of primary breast tumors (Nacht M, *et al.*, Cancer Research 59:5464-5470 (1999)). To confirm the validity of these findings, we investigated expression of cyclin D2 by RT-PCR in three finite life span and 6 immortal HMECs, 11 breast cancer cell lines and 24 primary breast carcinomas. A ribosomal protein RNA, 36B4, was co-amplified as an internal control. Abundant expression of cyclin D2 mRNA was noted in all three finite life span HMECs and in 4 of 6 immortalized HMECs. The two immortalized HMEC lines lacking cyclin D2 expression were HBL100 and MCF10A. In contrast, 10 of 11 breast cancer cell lines showed no detectable expression of cyclin D2. Only one breast cancer cell line, HS578T, expressed a low but detectable level of cyclin D2. Likewise, the results with primary tumors reflected the findings in cultured cells. Eighteen of 24 primary breast carcinomas expressed significantly lower levels of cyclin D2 mRNA as compared with finite lifespan HMEC 184 and five other HMECs. As an additional control for cyclin D2 expression, the expression of cyclin D1 was analyzed in the same panels of cell lines and tumors. Consistent with previous observations Cyclin D1 mRNA was detectable in all the cell lines and primary breast tumors tested. Thus, in both breast cancer cell lines and primary tumors specific loss of cyclin D2, but not cyclin D1, mRNA expression was observed.

[0118] Cyclin D2 mRNA expression in luminal and myoepithelial cells of the breast It has been reported that cyclin D2 is expressed in myoepithelial but not in luminal epithelial cells of the breast (Lukas J, (1995)). Therefore, lack of expression of cyclin D2 in breast cancers would be expected, since the vast majority of these tumors originate from luminal

rather than myoepithelial cells. This conclusion was based, however, on the results from a single HMEC preparation. The present study used a larger panel of tissues. Luminal and myoepithelial cells isolated from four normal mammaplasty specimens from women aged 18 to 33 were used. Paired luminal and myoepithelial cells were obtained from the same breast of two women. Each cell type was purified using immunomagnetic beads. The human luminal and myoepithelial cells were separated by virtue of their exclusive expression of epithelial membrane antigen (EMA) and common acute lymphoblastic leukemia antigen (CALLA) respectively. The purity of the populations was checked by immunocytochemistry using cytokeratins 18 and 19 as markers for luminal cells and cytokeratin 14 as a marker for myoepithelial cells. These tests showed that the final population was 95-99% pure in each case. Cyclin D2 expression was assessed in the purified cell preparations by RT-PCR. Cyclin D2 expression was observed in four of four purified luminal epithelial cells, as well as four of four myoepithelial cells. However, one luminal epithelial cell sample had a significantly lower expression of cyclin D2. Four HMECs of the 184 series, which stain for luminal cell markers- cytokeratins 8 and 18 and mucin, but not for myoepithelial cell marker- cytokeratin 14, also expressed cyclin D2 mRNA. Thus, cyclin D2 mRNA was expressed in all eight of eight luminal and four of four myoepithelial cell preparations from the normal breast.

[0119] Western analysis reveals loss of cyclin D2 protein in primary tumors For Western blot analysis specific anti-cyclin D2 antibodies that did not cross-react with cyclin D1 were used. While cyclin D2 protein was clearly detected in all seven HMECs tested (11-24, 1-26, 70N, 166372, 81N, 9F1403 and 184A1), it was undetectable in the majority (10/13) of primary breast tumors. Thus, HMECs that were derived from normal breast tissue and expressed high levels of cyclin D2 mRNA show clearly detectable levels of cyclin D2 protein as well. In contrast, primary breast tumors that exhibited low or absent cyclin D2 mRNA showed a corresponding loss of the cyclin D2 protein.

[0120] The cyclin D2 promoter is hypermethylated in breast cancer cell lines and primary tumors In somatic cells, about 80% of the CGs are methylated. Exceptions to this are the CpG islands in the promoter region of many genes. CpG islands are GC-rich regions

[0125] Expression of cyclin D2 protein was undetectable in 10 of the 13 primary breast cancers tested. However, methylation of the cyclin D2 promoter was noted only in six of

these ten primary tumors. This finding suggests that while methylation may cause silencing of cyclin D2 expression in many breast cancers, alternative pathways account for the loss of the protein in a proportion of these tumors.

[0126] Cyclin D2 promoter hypermethylation in preneoplasia Ductal carcinoma *in situ* (DCIS) is a preneoplastic lesion with a potential for progression to invasive cancer. To determine if hypermethylation of the cyclin D2 promoter occurs early in the evolution of breast cancer, MSP analysis was performed on DNA from carefully microdissected samples of DCIS. Hypermethylation was noted in 44% of DCIS samples. In the cases where adjacent invasive cancer was present as well, the methylation status of both lesions was concordant. This finding suggests that alteration of cyclin D2 expression may be an early event, and may precede transformation to the fully malignant stage of invasive carcinoma.

[0127] Re-expression of cyclin D2 mRNA in breast cancer cell lines Breast cancer cell lines MDAMB231 and MCF7 do not express cyclin D2 mRNA or protein. If silencing of expression is mediated by promoter methylation and/or altered chromatin conformation, then demethylation of the gene by exposure to 5-aza 2'-deoxycytidine (5aza-dC), or treatment with the histone deacetylase inhibitor, trichostatin A (TSA), should result in removal of the repressive mechanism and re-expression of the gene. Indeed, when MDAMB231 and MCF7 cells were exposed to 5-aza-dC in culture, the cyclin D2 promoter was partially demethylated (as analyzed by MSP), and cyclin D2 mRNA expression was restored (as analyzed by RT-PCR). Further, exposure to TSA also led to re-expression of the cyclin D2 mRNA. These results suggest that methylation at the promoter region plays a functional role in suppressing the expression of cyclin D2 in breast cancer.

[0128] Using RT-PCR, cyclin D2 expression was detected in four normal luminal epithelial cultures of the 184 series, in four of four purified luminal epithelial cell extracts, and in four of four myoepithelial cell extracts. Using MSP, promoter hypermethylation was detected in 49/106 (46%) of the tumors. Hypermethylation of the gene correlated with lack of cyclin D2 mRNA and/or protein expression. Thus, in about 50% of breast cancers, cyclin D2 silencing may be attributed to tumor-specific methylation.

EXAMPLE 3

Hypermethylation and loss of expression of 14-3-3 sigma

[0129] The extent of methylation of the cyclin 14-3-3 sigma-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors, and expression of the 14.3.3 sigma mRNA and protein in the same cells and tissues was examined.

[0130] **Cell Lines and Tissues** The breast cancer cell lines Hs578t, MDA-MB-231, MDA-MB-435 and MCF-7 and the human mammary epithelial cell lines, MCF-10A and HBL-100 were obtained and maintained according to instructions (ATCC). The two matched tumor cell lines, 21PT and 21MT were propagated as described (Band, *et al.* (1990) *Cancer Res.* 50:7351-7357). Cultured normal human breast epithelial cell (HMEC) strains, 161, 184, 172, and 48, and the conditionally and fully immortal cell lines, 184A1 (passage 15 and 99), and 185B5 were grown as described (<http://www.lbl.gov/LBL-Programs/mrgs/review.html>). Three additional short term cultures of HMECs, (#04372 and #16637) were grown according to specifications (Clonetics). Primary breast tumor tissues were obtained immediately after surgical resection at the Johns Hopkins University or Duke University, and stored frozen at -80°C. Microscopic examination of representative tissue sections from each tumor revealed that these samples contained greater than 50% tumor cells. Microdissection of primary tumor cryosections was performed by using a laser capture microscope (Schutze, *et al.* (1998) *Nat Biotechnol* 16:737-42) or by manually scraping the cells with a 20G needle under 40X magnification (Umbricht, *et al.* (1999) *Oncogene* 18:3407-14.).

[0131] **Northern Blot Analysis** Total RNA was isolated from primary tumor tissues using Trizol Reagent (Life Technologies). Five micrograms were resolved on 1.5% agarose/formaldehyde gels, and transferred to a nylon filter using standard methods (Gene Screen, DuPont). A 375 bp σ -specific probe was generated using MCF-10A cDNA as a template and the primers 5'-ACAGGGGAAC TTTATTGAGAGG-3' (SEQ ID NO:25) and 5'-AAGGGCTCCGTGGAGAGGG-3' (SEQ ID NO:26). Hybridizations were done in Quikhyb (Stratagene) according to the manufacturer's instructions. Filters were exposed to

autoradiographic film for up to 5 days. To test for uniform loading of the samples, blots were stripped and reprobed with a 1.5 kb DNA fragment specific for 18S rRNA (ATCC, Clone #HHCSA65).

[0132] Loss of Heterozygosity (LOH) Studies A TG repeat sequence in the 3'UTR of σ was amplified using: 5'-GAGGAGTGTCCTGCTGG-3' (sense; SEQ ID NO:27) and 5'-GTCTCGGTCTTGCACTGGC-3' (antisense; SEQ ID NO:28) primers, which yields a product of 117 bp. The 25 μ l reactions contained 50 ng of template DNA (10), 17 mM NH₄SO₄, 67 mM TrisCl (pH 8.8), 6.7 mM MgCl₂, 1% DMSO, 1.5 mM dNTP, 20 ng of each primer, 2 ng of γ -³²P-labeled sense primer, and 0.5 μ l Taq polymerase. PCR conditions were as follows: 1 cycle of 94 °C for 90s; 35 cycles of 94 °C for 1 min, 57 °C for 30s, 72 °C for 30s; and 1 cycle of 72 °C for 5 min. PCR products were fractionated on a sequencing gel, which was exposed to autoradiographic film overnight (Evron, *et al.* (1997) *Cancer Res*, 57:2888-9).

[0133] Mutation Analysis A 1.2 kb PCR product, encompassing the entire σ coding sequence, was generated using two primers, 5'-GTGTGTCCCCAGAGCCATGG-3' (sense; SEQ ID NO:29) and 5'-GTCTCGGTCTTGCACTGGCG-3' (antisense; SEQ ID NO:30). The PCR reaction contained 50 ng of DNA, 6.4% DMSO, 1.5 mM dNTPs, 100 ng of each primer and 0.5 μ l Taq polymerase in a 50 μ l reaction volume. α -³³P cycle sequencing was performed using the Amplicycle sequencing kit (Perkin Elmer). Four different α -³³P-labeled primers were used to sequence the entire σ coding sequence:
5'-CACCTTCTCCCGGTACTCACG-3' (antisense; SEQ ID NO:31),
5'-GAGCTCTCCTGCGAAGAG-3' (sense; SEQ ID NO:32),
5'-GAGGAGGCCATCCTCTCTGGC-3' (sense; SEQ ID NO:33) and
5'-TCCACAGTGTGAGGTTGTCTCG-3' (antisense; SEQ ID NO:34).

[0134] Transfection of Human Breast Cancer Cell Lines 1.5 x 10⁵ of MCF-7, MDA-MB-231, and Hs578t, or 2.5 x 10⁵ cells of MDA-MB-435 breast cancer cells were seeded in six-well plates. The following day, transfections were performed using Trans IT-LT1 (Mirus Corp.) as per manufacturer's instructions. Plasmids used in the transient

transfections include: KKH luciferase, containing 4 kb of the σ -promoter linked to the luciferase gene in the pGL3-Basic vector (Promega); pCMV- β -gal (Clontech), which was used to correct for the efficiency of transfection; and pGL3-Basic (Promega), which was used as a negative vector control against which KKH luciferase activities were compared. Two μ g of luciferase reporter plasmid or the pGL3-Basic vector control and 0.5 μ g of CMV- β -gal reporter plasmid were used for each transfection.

[0135] Luciferase and β -galactosidase Assays Cell lysates were made approximately 48 hr post-transfection as per manufacturer's instructions (Promega, Luciferase Assay System). Luciferase and β -galactosidase activities were quantitated using the luciferase assay system (Promega) and the Aurora GAL-XE® reporter gene assay (ICN Pharmaceuticals, Inc), respectively. Experiments were done in triplicate. Luciferase activity was first normalized for efficiency of transfection by using the ratio of luciferase to β -galactosidase activity. For each transfected cell line, the results were compared with the mean of pGL3 vector control levels and expressed as fold elevated expression above pGL3. The means and standard deviations of the results of all experiments were calculated.

[0136] Sodium Bisulfite DNA Sequencing Genomic DNA was subjected to sodium bisulfite modification as described in Herman *et al.* ((1996) *Proc. Natl. Acad. Sci. USA*, 93:9821-9826). Bisulfite-converted DNA was amplified, as described above, using primers that encompass the first exon of the σ gene: 5'-GAGAGAGTTAGTTTGATTAGAAAG-3' (sense primer with start at nt 8641; SEQ ID NO:35) and 5'-CTT ACTAATATCCATAACCTCC-3' (antisense primer with start at nt 9114; SEQ ID NO:36) which generated a 474 bp PCR product. Conditions for PCR were as follows: 1 cycle at 95 °C for 5 min; 35 cycles at 95 °C for 45s, 55 °C for 45s and 72 °C for 60s; and 1 cycle at 72 °C for 4 min. The product was purified using a Qiagen PCR purification kit (Qiagen Corp) and sequenced using the sense primer with an ABI automated fluorescent sequencer according to the manufacturer's instructions.

[0137] Methylation-specific PCR (MSP) One μ g genomic DNA was treated with sodium bisulfite as described in (Herman, *et al.*. (1996) *Proc. Natl. Acad. Sci. USA* 93,

9821-9826), and was analyzed by MSP using a primer set that covered CG dinucleotide numbers 3, 4, 8 and 9. Primers specific for methylated DNA: 5'-TGGTAGTTTTTATGAAAGGCGTC-3' (sense; SEQ ID NO:37) and 5'-CCTCTAACCGCCCACCACG-3' (antisense; SEQ ID NO:38), and primers specific for unmethylated DNA: 5'-ATGGTAGTTTTTATGAAAGGTGTT-3' (sense; SEQ ID NO:39) and 5'-CCCTCTAACCACCCACCACA-3' (antisense; SEQ ID NO:40) yielded a 105-107 bp PCR product. The PCR conditions were as follows: 1 cycle of 95 °C for 5 min; 31 cycles of 95 °C for 45s, 56 °C for 30s and 72 °C for 30s; and 1 cycle of 72 °C for 4 min.

[0138] Treatment of Cells with 5'-aza-2'-deoxycytidine (5-aza-dC) Cells were seeded at a density of 2×10^6 cells per 100-mm plate. 24 h later cells were treated with 0.75 μ M 5-aza-dC (Sigma) (Ferguson, *et al.* (1995) *Cancer Res* 55:2279-2283). Total cellular RNA and genomic DNA were isolated from the cells at 0 and 3 days after addition of 5-aza-dC as described herein.

[0139] RT-PCR RNA was treated with RNase-free DNase (Boehringer-Mannheim) (1 μ g/ μ l) for 2h at 37° C, followed by heat inactivation at 65 °C for 10 min. RT reactions contained 1 μ g DNase treated RNA, 0.25 μ g/ μ l pdN6 random primers (Pharmacia), 1x first strand buffer (GibcoBRL), 0.5 mM dNTP (Pharmacia), and 200 U MMLV-RT (GibcoBRL), and were incubated for 1h at 37° C. PCR was performed using the σ -specific primers 5'-GTGTGTCCCCAGAGCCATGG-3' (SEQ ID NO:41) and 5'-ACCTTCTCCCGGTACTCACG-3' (SEQ ID NO:42) using buffer conditions described herein. The PCR conditions were: 1 cycle of 95° C for 5 min; 30 cycles of 60° C for 45s, 72° C for 45s and 95° C for 45s. PCR samples were resolved by electrophoresis in a 2% agarose gel.

[0140] Assay for G1 and G2 checkpoint and chromosomal aberrations The G1 cell cycle checkpoint and chromosomal aberrations in mitosis were assessed as described previously (Pandita, *et al.* (1996) *Oncogene* 13:1423-1430). Specifically, cells in plateau phase were irradiated with 3 Gy, sub-cultured after 24h, and metaphases were collected. G1 type aberrations were examined at metaphase. All categories of asymmetric chromosome

105579-01802

aberrations were scored: dicentrics, centric rings, interstitial deletions/acentric rings, and terminal deletions.

[0141] The efficiency of G2 checkpoint control was evaluated by measuring the proportion of cells in metaphase after irradiation. Chromosomal aberrations at mitosis were assessed by counting chromatid breaks and gaps per metaphase as described elsewhere (Morgan, *et al.* (1997) *Mol Cell Biol* 17:2020-2029). Specifically, cells in exponential growth phase were irradiated with 1 Gy. Metaphases were harvested 45 and 90 minutes following irradiation and examined for chromatid type breaks and gaps. Fifty metaphases each were scored for G1 and G2 types of chromosomal aberrations.

[0142] **Introduction of σ into the σ -negative breast cancer cell line MDA-MB-435 by adenoviral infection** Cells were seeded and grown to 50% confluence. Adenovirus encoding either σ or β -galactosidase (Hermeking, *et al.* (1997) *Mol. Cell* 1:3-11) was added to the culture at a multiplicity of infection of 5000:1 and infection was allowed to take place overnight. The cells were harvested, fixed and stained with Hoechst dye and subjected to FACS analysis.

[0143] **σ Expression in Normal, Immortalized and Tumorigenic Breast Epithelial Cells** By SAGE analysis, σ was found to be expressed at an average of 7-fold lower levels in three human breast cancer cell lines, 21PT, 21MT and MDA-MB-468 than in two populations of normal human mammary epithelial cells (HMEC). Northern blot analysis was performed to confirm this finding in other breast cancer cell lines and in primary breast tumors. No expression of σ was detected in 45 of 48 (94%) primary tumors. In contrast, σ was expressed at easily detectable levels in all 6 cultured HMEC populations and 5 immortalized but nontumorigenic cell lines. These results indicate that loss of σ gene expression is a frequent event in human breast cancer.

[0144] **Genetic alterations within the σ gene** Possible causes for loss of σ gene expression in breast tumors include deletion of the chromosomal region containing the gene or intragenic mutations that lead to decreased mRNA stability. σ localizes to chromosome 1p35, an arm that has been extensively studied for LOH in breast cancer (Hermeking, *et al.*

SECRET

(1997) *Mol. Cell* **1**, 3-Bieche, *et al.* (1995) *Genes Chrom. Cancer* **14**:227-251). LOH has been observed for the 1p32-36 region at a frequency of 15-25%. However, it is not known whether the region lost in these tumors includes σ (Genuardi, *et al.* (1989) *Am. J. Hum. Genet.* **45**:73-82; Trent, *et al.* (1993) *Genes Chrom Cancer* **7**:194-203; Nagai, *et al.* (1995) *Cancer Res.* **55**:1752-1757; Tsukamoto, *et al.* (1998) *Cancer* **82**:317-322). Therefore, the loss of σ by utilizing a TG repeat sequence within the 3' UTR of the σ gene itself was examined. Using primers that span the TG repeats, the locus in 45 sets of normal and tumor DNA pairs was studied. Twenty of 45 (44%) of the patients were found to be heterozygous with respect to the length of the PCR-product. Only one of the 20 tumor specimens exhibited LOH (Table 2). Eleven of these 20 samples were tested by Northern blot analysis, and no σ transcripts were detectable. These results prompted an examination whether there were smaller genetic changes within the coding region of σ . The entire 1190 bp coding region from σ -nonexpressing (σ -negative) breast cancer cell lines, MDA-MB-435 and Hs578t and 7 primary tumor tissues was amplified with PCR and sequenced. No mutations were found. In addition, 25 primary tumor DNA samples were analyzed by single stranded conformation polymorphism, and no abnormalities were detected. These results suggest that genetic alterations within σ are not a primary mechanism for loss of gene expression.

2025 RELEASE UNDER E.O. 14176

Table 2. Incidence of σ alterations in breast cancer

Sample	σ expression, Northern blot analysis	No. with methylated $\sigma\sigma$ /total		No. with LOH/total,	No. with mutation/total
		Sequencing	MSP	TG repeat PCR	Sequencing SSCP
Normal breast					
Mortal HMEC strains	6/6	01	0/3		
Immortal HMEC lines	5/5	01	0/5		
Reduction mammaplasty, microdissected epithelium			0/6		01
Breast cancer Cell Lines					
MCF-7	+	♂	♂		
MDA-MB-231	+	♂	♂		
MDA-MB-435	♂	+	+		♂
Hs578t	♂	+	+		♂
Primary tumors	2/45	10/10	43/50	1/20	0/7 0/25
MICRODISSECTED CARINOMA			32/32		

[0145] **Epigenetic alterations of the σ gene** Next tested was whether the lack of σ mRNA was due to deficiencies in factors required for σ transcription. The two breast cancer cell lines, MDA-MB-435 and Hs578t, served as model systems for σ -negative primary tumors that harbored wild type σ alleles, while the two breast cancer cell lines, MCF-7 and MDA-MB-231, served as σ -positive controls since they both express detectable levels of σ . The plasmid KKH-luciferase contains 4 kb of sequence upstream of the transcriptional start site of σ linked to the luciferase reporter gene; this upstream region contains the sequences necessary for p53 and γ -irradiation-inducible transcription of σ (5). Following transient transfection of the four cell lines with the reporter plasmid, high levels of expression was observed (70- to 300-fold above the promoterless parental vector) in both σ -negative and σ -positive breast cancer cell lines. These results indicate that the σ -negative breast cancer cells, like the σ -positive cells, are able to support transcription from the σ promoter equally well, and contained factors required for transcription.

[0146] σ has a CpG rich region (CpG island) within its first and only exon that begins near the transcription initiation site and ends approximately 800 bp downstream. To explore a role of hypermethylation in silencing σ gene expression, the nucleotide sequence of this region was determined after treating the DNA with sodium bisulfite (Frommer, *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:1827-1831). PCR primers were designed to amplify a region spanning 27 CpG dinucleotides within the CpG island. No significant methylation was observed using DNAs from four σ -positive cell lines including 2 HMECs (184, MCF-10A) and two tumorigenic breast cancer cell lines (MCF-7 and MDA-MB-231). In contrast, DNAs from two σ -negative breast cancer cell lines, HS578t and MDA-MB-435, were fully methylated at all of the CpG sites. Since there was a strong correlation between σ -methylation status and mRNA expression in all the cell lines examined, 10 σ -negative primary breast tumors were also examined. All of the tumor DNAs exhibited partial or complete methylation of the 27 CpG dinucleotides.

[0147] Next, an MSP assay was utilized to detect methylation of the CpG island, using primers spanning the region between CpG dinucleotides 3 and 9 within the σ gene. Primers

105570.01200
2003-10-06 15:00

were designed that take advantage of the nucleotide sequence differences between methylated and unmethylated DNA as a result of bisulfite modification. By this method, 5/5 σ -positive HMEC strains were completely unmethylated. In addition, DNAs from the σ -positive immortalized breast epithelial cells (MCF-10A, HBL-100) and breast cancer cell lines (MCF-7 and MDA-MB-231), were also unmethylated at the sites examined. In contrast, DNAs from the σ -negative breast cancer cell lines, Hs578t and MDA-MB-435, were fully methylated. Similarly, 43 of 50 samples from primary breast tumors were partially or completely methylated. Of these 43 tumors, 26 were examined by Northern blot analysis, and all 26 lacked detectable σ gene expression. Three of the seven unmethylated breast tumor samples also lacked σ transcripts; the expression pattern for the remainder was not tested. These results demonstrate that aberrant methylation of σ is a frequent event in breast cancer, but that other mechanisms are responsible for silencing the gene in a small fraction of breast tumors.

[0148] Previous reports indicate that σ gene expression is restricted to differentiated epithelial cells. In order to clearly ascertain the cellular origin of methylated DNA, normal and tumor tissues were microdissected and analyzed for σ methylation by MSP. All six DNA samples of microdissected mammary epithelial cells obtained from reduction mammaplasty specimens were unmethylated. In contrast, all 32 samples of DNA from microdissected breast carcinomas were methylated within the σ CpG island. These results indicate that hypermethylation of the σ gene is associated with loss of gene expression in the majority of primary breast tumors. The data from gene expression, genetic and epigenetic studies are summarized in Table2.

[0149] In order to determine the effect of methylation on σ gene expression, two fully methylated, σ -negative cell lines, Hs578t and MDA-MB-435, were treated with the DNA methyltransferase inhibitor, 5-aza-dC. Treatment of cells with 0.75 μ M 5-aza-dC for 3 days led to demethylation of the CpG rich region encompassed by the MSP primers. Moreover, 5-aza-dC treatment resulted in reactivation of gene expression, as demonstrated by RT-PCR. These results demonstrate that methylation is at least partially responsible for loss of σ transcription in breast cancer cells.

[0150] Functional consequences of loss of σ in breast cancer cells The function of human σ has been analyzed in human colon carcinoma cells. These studies demonstrated that following ionizing irradiation, σ sequesters cdc2-cyclin B1 complexes in the cytoplasm, thus arresting the cells in G2. These actions prevent the cell from initiating mitosis before repair of its damaged DNA. Colon carcinoma cells lacking σ can still initiate, but do not maintain, G2 arrest, leading to mitotic catastrophe and cell death.

[0151] In an attempt to determine the effects of loss of σ gene expression on cell cycle regulation in breast cancer cells, the effects of γ -irradiation on the σ -negative breast cancer cell lines, MDA-MB-435, 21NT, and 21MT, and the σ -positive breast cancer cell line, MCF-7 were tested. First, G1 type chromosomal aberrations were examined 24 h after cells were exposed to 3 Gy of γ -irradiation. All categories of G1-type chromosomal aberrations were scored at metaphase; their frequency was identical in the two cell types. These results indicate that the examined cell lines have similar G1 cell cycle checkpoint control responses to ionizing radiation.

[0152] Next, the G2 checkpoint function in the four cell lines was evaluated. Cells in exponential growth phase were γ -irradiated with 1 Gy and metaphases were examined for chromatid type breaks and gaps. Defective G2 arrest will increase these values. The results show a striking difference in the ability of σ -negative cells and σ -positive cells to repair their damaged DNA. Forty-five minutes post irradiation, σ -negative cells exhibited up to twice as many G2 type chromosomal aberrations as MCF-7 cells. This number increases to three-fold by 90 minutes. Moreover, while repair of DNA damage was evident in the MCF-7 cells, as evidenced by a decrease in the number of G2 type aberrations between 45 and 90 minutes, no decrease was seen in σ -negative cells.

[0153] Finally, in order to further demonstrate the role of σ in G2 checkpoint function in breast cells, a cloned copy of the gene was overexpressed in the σ -negative cell line MDA-MB-435 as well as in normal breast epithelial cells using the adenovirus expression system used to express σ in colon cancer cells (5). Overexpression of σ in these breast epithelial cells led to a rapid and permanent G2 arrest, whereas the control virus-infected

1005079.01202

cells showed no effect. These results indicate that although the σ -negative cell lines have a functional G1 cell cycle checkpoint, they accumulate more genetic damage following irradiation, which is consistent with its failure to arrest in G2 in response to DNA damage.

[0154] In summary, these results show that in striking contrast to normal breast tissue, greater than 90% of breast cancers lack detectable expression of σ . Hypermethylation of the σ gene occurs in a CpG-rich region that extends from the transcriptional initiation site to the middle of the coding region. Bisulfite genomic sequencing of this 500 bp region showed that is consistently and densely methylated in σ -negative cell lines and primary breast tumors. Several studies have clearly documented that gene activity correlates inversely with the density of gene-specific CpG island methylation, but is less dependent on the position and distance of the methylated DNA sequences from the transcriptional initiation site. With respect to σ , dense methylation just downstream of its transcriptional start site is strongly associated with gene silencing. Furthermore, in σ -negative cell lines, 5-aza-dC-induced demethylation of the CpG island leads to reactivation of gene expression, indicating that hypermethylation plays a causal role in σ gene inactivity.

EXAMPLE 4

Hypermethylation and loss of expression of RAR β 2

[0155] The extent of methylation of the RAR β 2-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors, and expression of the RAR β 2 mRNA and protein in the same cells and tissues was examined.

[0156] **Cell cultures** Human epithelial mammary cells (HEMC) from reduction mammaplasty including three mortal strains, 184, 48R and 172R, and two immortal strains, 184A1 and 184B5, were obtained and cultured according to the protocols designed by Dr. Martha Stampfer (see the HMEC Homepage, http://www.lbl.gov/*mrgs/index.html) using Clonetics (Walkersville, MD, USA) reagents. Human breast cancer cell lines were maintained in Dulbecco's modified Eagle's medium (GIBCO) (Hs578t, MCF-7, MDA-MB-231 and T47D) or IMEM medium (Biofluids) (MDA-MB-435, MDA-MB-468,

ZR751) with 5% fetal calf serum (FCS). For drug treatments, exponentially growing cells were seeded in 10 cm² plates at a density of 36105 cells/plate or in 6-well plates at 16105 cells/well. Cells were allowed to attach overnight before the addition of the appropriate concentration of 5-Aza-2' deoxycytidine (5-Aza-CdR) (Sigma), Trichostatin A (TSA) (Sigma) or RA(Sigma). When reduction of retinoids was required, cells were treated in either medium with 0.5% FCS or charcoal-dextran stripped FCS (Hyclone). At the indicated time points, both attached and detached cells were harvested, counted with Trypan Blue (Life Technologies) and processed for DNA or RNA extraction. 5-Aza-CdR was dissolved in 0.45% NaCl containing 10 mM sodium phosphate (pH 6.8). Trichostatin A and all-trans-retinoic acid (RA) (Sigma) were reconstituted in absolute ethanol (solvent). The growth inhibition (%) was calculated as: $(1 - NT/NC) \times 100$, where NT is the number of treated cells and NC is the number of control cells.

[0157] Tissue samples Normal and tumor tissues were collected from existing tumor banks (Istituto per lo Studio e la Cura dei Tumori, Milan; the Cancer Center, Rotterdam, the Johns Hopkins Breast Cancer Program, Baltimore, MD, USA). All tumor samples were obtained from excess clinical specimens and institutional guidelines for the acquisition and maintenance of such specimens were followed. DNA and RNA extraction: Extraction of DNA and RNA from breast cancer cell lines was performed by using DNAzol and Trizol respectively (LifeTechnologies) according to the manufacturer's instructions. Genomic DNA was further treated with 500 mg/ml proteinase K at 55°C, extracted with phenol-chloroform-isoamyl alcohol (24 : 24 : 1) (CIA) and ethanol precipitated. Extraction of DNA from paraffinated breast cancer and lymph node tissues was essentially performed as previously described (Formantici et al., 1999). One to three consecutive sections estimated to contain at least 90% tumor cells were incubated at 58°C overnight in 200 ml of extraction buffer (50 mM KCl, 10 mM Tris-HCl (pH 7.5), 2.5 mM MgCl₂, 0.1 mg/ml gelatin, 0.45% NP-40, 0.45% Tween 20, and the solution was heated at 95°C for 15 min to inactivate the proteinase K and then centrifuged at 6000 r.p.m. The DNA in the supernatant was used for analysis.

[0158] Southern blotting Genomic DNA (7 mg) was digested overnight with 15 U/ mg of XbaI, HpaII and MspI enzymes, electrophoresis on a 0.8% agarose gel and transferred to

Hybond-N filter. A 227 bp probe was amplified using the sense 5'-AGA GTT TGA TGG AGTTGG GTG GAG-3' (SEQ ID NO:43) and antisense 5'-CAT TCG GTT TGGGTC AAT CCA CTG-3' (SEQ ID NO:44) primers, gel purified and labeled with ³²P-dCTP using the Megaprime DNA labeling system (Amersham). After hybridization the filters were washed and exposed to X-ray film at -80°C for autoradiography.

[0159] Methylation specific PCR (MSP) Bisulfite modification of genomic DNA was essentially performed as described by Herman et al. (1996) and described herein. Modified DNA was used immediately or stored in aliquots at -20°C. The PCR mixture contained 1 x PCR buffer (16.6 mM ammonium sulfate, 67 mM Tris (pH 8.7), 1.5 mM MgCl₂), dNTPs (each at 1.25 mM), primers (300 ng each per reaction), and bisulfite-modified DNA (50 ng) or unmodified DNA (50 ng). Reactions were hot started at 95°C before the addition of 2.5 U of Taq polymerase (Qiagen). Amplification was carried out in a Thermal Cycler 480 Perkin Elmer for 30 cycles (1 min at 94°C, 1 min at the annealing temperature (at) selected for each primer pair, 1 min at 72°C), followed by 4 min at 72°C. Twelve µl of the PCR reaction were electrophoresed onto 1.5% agarose gels, stained with ethidium bromide and visualized under UV. Two primer pairs, W3 sense 5'-CAGCCCGGGTAGGGTTCACC-3' (SEQ ID NO:45), W3 antisense 5'-CCGGATCCTACCCCGACGG-3' (SEQ ID NO:46), and W4 sense 5'-CCGAGAACGCGAGCGATCC-3' (SEQ ID NO:47) and W4 anti-sense 5'-GGCCAATCCAGCCGGGCG-3' (SEQ ID NO:48), were designed on the human RAR β2 sequence (Shen *et al.*, 1991) and used to control the Na bisulfite modification. The primer pairs selected to detect the unmethylated DNA were as follows: U1 sense 5'-GTG GGT GTA GGT GGA ATA TT-3' (SEQ ID NO:49) and U1 antisense 5'-AAC AAA CAC ACA AAC CAA CA-3' (SEQ ID NO:50) (at 55°C); U2 sense 5'-TGT GAG TTA GGA GTA GTG TTTT-3' (SEQ ID NO:51) and U2 antisense 5'-TTC AAT AAA CCC TAC CCA-3' (SEQ ID NO:52) (at 49°C); U3 sense 5'-TTA GTA GTT TGG GTA GGGTTT ATT-3' (SEQ ID NO:53) and U3 antisense 5'-CCA AAT CCT ACC CCAACA-3' (SEQ ID NO:54) (at 55°C); U4 sense 5'-GAT GTT GAG AAT GTGAGT GAT TT-3' (SEQ ID NO:55) and U4 antisense 5'-AAC CAA TCC AACCAA AAC A-3' (SEQ ID NO:56) (at 55°C); The sequences of the primers to detect the methylated DNA were: M1 sense 5'-AGC GGGCGT AGG CGG AAT ATC-3' (SEQ ID NO:57) and M1 antisense 5'-CAACGA ACG CAC AAA CCG ACG-3'

SEQUENCE LISTING

(SEQ ID NO:58) (at 63°C); M2 sense 5'-CGT GAG TTA GGA GTA GCG TTT C-3' (SEQ ID NO:59) and M2 antisense 5'-CTT TCG ATA AAC CCT ACC CG-3' (SEQ ID NO:60) (at 57°C); M3 sense 5'-GGT TAG TAG TTC GGG TAG GGT TTA TC-3' (SEQ ID NO:61) and M3 antisense 5'-CCG AAT CCT ACC CCGACG-3' (SEQ ID NO:62) (at 64°C); M4 sense 5'-GTC GAG AAC GCG AGCGAT TC-3' (SEQ ID NO:63) and M4 antisense 5'-CGA CCA ATC CAA CCGAAA CG-3' (SEQ ID NO:64) (at 64°C).

[0160] M and U primers were designed in the same regions, with one or two nucleotide differences to meet annealing requirements. Fragment M3 (position 773 ± 1007) contains the βRARE (792 ± 808) and the transcription start site (position 844); fragment M4 (position 949 ± 1096) contains an Sp1 element (position 1074 ± 1081).

[0161] **RT± PCR** The exon 5 (sense primer 5'-GAC TGT ATG GAT GTTCTG TCA G-3'; SEQ ID NO:65) and exon 6 (antisense primer 5'-ATT TGTCCT GGC AGA CGA AGC A-3'; SEQ ID NO:66) were designed on the basis of published RAR β2 transcript (de The' et al., 1990; van der Leede et al., 1992) and used to amplify 50 ng of DNase treated total RNA using the Superscript One-StepRT± PCR System (Life Technologies). RT± PCR with actin primers (sense primer 5'-ACC ATG GAT GAT GAT ATCG-3'; SEQ ID NO:67 and antisense primer 5'-ACA TGG CTG GGG TGTTGA AG-3'; SEQ ID NO:68) was used as an internal RNA control.

[0162] **The RAR β2 promoter is methylated in breast cancer cell lines independently of their ER status and RA-inducibility** RAR transcription was first tested in a panel of breast cancer cell lines grown in the absence of exogenous RA, by reverse transcriptase-PCR (RT ± PCR), using primers encompassing exons 5 and 6 (de The' et al., 1990; van der Leede et al., 1992; Toulouse et al., 1997). Under these conditions, only one cell line, Hs578t, produced a detectable 256 bp RT± PCR product. Thus, previous reports were confirmed that RAR β gene expression is down regulated/lost in breast cancer cell lines. Growing cells in the presence of RA can assess the distinction between down regulation and loss. As previously reported (Swisshelm et al., 1994; Liu et al., 1997; Shang et al., 1999), we observed induction of RAR β expression and growth inhibition in T47D, MDA-MB-435,

1005579-013802

MCF7 and ZR75-1 cell lines treated for 48 h with 1 μ M RA, but not in the MDA-MB-231 and MDA-MB-468 cell lines.

[0163] To see whether the RAR β 2 methylation status correlated with the ER status, the methylation status was examined at RAR β 2 in a panel of ER-positive (MCF7, T47D, ZR75-1) and ER-negative (Hs578t, MDA-MB-231, MDA-MB-435, MDA-MB-468) cell lines.

[0164] By Southern blotting, the CpG island of the RAR β 2 promoter within a 7.5 kb XbaI DNA fragment encompassing the TATA box, the β RARE, the transcriptional start site (TS) and the 5' untranslated region of exon 5 was examined. In this region nine HpaII sites can be identified (Shen *et al.*, 1991; Baust *et al.*, 1996). The DNA methylation status was analyzed by using the methylation-sensitive enzyme, HpaII. MspI, the isoschizomer of HpaII, insensitive to methylation, was used as a positive control. The PCR probe spans the β RARE and the TATA box regions. The same 7.5 kb region was previously analyzed in a colon carcinoma cell line, and the size of all the possible fragments relative to the most 3' HpaII site were reported (Cote' and Momparler, 1997). Genomic DNA from the ER-positive, RA-inducible cell line T47D is digested to completion, indicating that it is not methylated at any of the HpaII sites. In contrast, DNA from the ER-positive, RA-inducible ZR75-1 cell line and DNA from the ER-negative, RA-resistant MDA-MB-231 cell line showed to be differentially methylated at the methylation-sensitive sites. Using methylation-specific PCR (MSP), we further analyzed a 616 bp long RAR β 2 region from nucleotide 481 to nucleotide 1096 (Shen *et al.*, 1991) in all the cell lines. MSP entails the modification of genomic DNA by sodium bisulfite that converts all unmethylated, but not methylated, cytosine to uracil (Herman *et al.*, 1996). The genomic DNAs from four breast cancer cell lines ZR751, MCF7, MDA-MB-231, MDA-MB-468 showed partial to complete methylation of the promoter region. The human mammary epithelial cell (HMEC) strain 48R, expressing RAR β and three breast cancer cell lines, the RAR β -positive Hs578t and the RA-inducible MDA-MB-435 and T47D, revealed only the (U) unmethylated PCR products.

10065570.01.0000

[0165] These results indicate that hypermethylation of the RAR β 2 promoter occurs in breast cancer cell lines irrespective of the ER status, and can be detected in both RA-inducible, and RA-resistant breast cancer cells.

[0166] **RAR β 2 is unmethylated in both mortal and immortalized HMEC, but is methylated in primary breast tumors** The next question examined was whether hypermethylation of RAR β 2 promoter in cell lines has correlates in clinical breast cancer. As a normal control, the HMEC mortal strains (48R, 172R), that are the closest representation of normal mammary epithelial cells available were examined. Also analyzed were two immortal mammary epithelial strains (184A1 and 184B5). The DNA of these strains was found to be unmethylated. Consequently, methylation of RAR β 2 may be an event in the progression of breast cancer, following immortalization. Genomic DNAs from three paraffinated samples of breast tumors, two ER-positive (T1, T2) and one ER-negative (T3), estimated to contain more than 90% tumor cells, were analyzed with all MSP primer pairs, and shown to be partially methylated. Both microdissected breast stroma, and microdissected normal epithelial cells were found unmethylated at RAR β 2, making it very likely that the U products in the tumor samples were amplified either from residual normal epithelial cells, or stromal cells mixed to tumor cells. DNAs from matching histologically tumor free lymph node samples ($N1 \pm N3$), were similarly analyzed and produced only the unmethylated PCR products. The DNA of additional 21 tumors was performed using two sets of primer pairs (U3/M3 and U4/M4). Fifteen (7 ER-positive and 8 ER-negative) of the 24 tumors presented methylation at the RAR β 2 promoter. With the same primer sets hypermethylation at RAR β 2 was detected in the DNA of ten out of 39 primary breast tumors collected, and analyzed independently, at the Johns Hopkins University. The overall data indicate that hypermethylation at RAR β 2 promoter occurs in approximately one third of primary breast tumors, and that the RAR β 2 methylation state is independent of the ER status of the tumor.

[0167] **5-Aza-CdR induces partial demethylation at the RAR β 2 CpG island and reactivation of RAR β gene expression** In order to determine whether DNA methylation is affecting, at least in part, RAR β gene expression, all cell lines showing methylation at the

10059579.016602

RAR β promoter were treated with the DNA methyl transferase inhibitor, 5-Aza-CdR. Treatment of cells with either 0.4 or 0.8 mM 5-Aza-CdR for 3 days, led to partial demethylation of the CpG rich RAR β region. This was evident both by Southern analysis in the MDA-MB-231 cell line, and by MSP in all cell lines. Moreover, 5-Aza-CdR treatment resulted in reactivation of gene expression both in RA-inducible MCF7 and ZR75-1, and RA-resistant MDA-MB-231 and MDA-MB-468 cells. Subsequent studies examined whether reactivation of RAR β expression by 5-Aza-CdR-resistant cells could be enhanced by RA. Using non-quantitative RT-PCR, a difference could not be appreciated in the level of RAR β transcription in MDA-MB-231 cells treated with 0.4 mM 5-Aza-CdR alone, or in combination, with 1 μ M RA. In this experiment, 5-Aza-CdR alone, or in combination with RA, produced 63 and 96% growth inhibition respectively. In the same experiment, treatment with 1 μ M RA alone produced a negligible effect on growth inhibition (52%). A synergistic effect of the two drugs on cancer cells was previously reported (Cote' and Momparler, 1997; Bovenzi *et al.*, 1999).

[0168] These data indicate that DNA methylation is, at least, one factor influencing the down regulation/loss of RAR β transcription in breast cancer cell lines with a methylated RAR β promoter. Cells treated with 5-Aza-CdR alone, or in combination with RA, showed re-expression of RAR β , which may have contributed, along with the toxic 5-Aza-CdR, to the observed growth inhibition.

[0169] **The HDAC inhibitor TSA can reactivate RAR β expression in RA-resistant cells; demethylation of the RAR β promoter is not an absolute requirement for RAR β reactivation** The chromatin status at a given locus can be dynamically influenced by the degree of acetylation/deacetylation due to HAT/HDAC activities. Absence of RAR β regulatory factors, like RAR α , as well as DNA-methylation, can contribute to pattern chromatin modifications at RAR β promoter in RA-resistant cell lines. One of these cell lines, MDA-MB-231, lacks RA-inducible RAR α activity (Shao *et al.*, 1994) and displays a RAR β methylated promoter. A subsequent study was designed to probe indirectly whether the level of HDAC at RAR β can influence RAR β expression, by testing the effect of TSA, a HDAC inhibitor on MDA-MB-231 cells (Yoshida *et al.*, 1995). Cells were treated for 2

days, in the presence or absence of 100 ng/ml TSA alone, or in combination, with 1 μ M RA. By using RT \pm PCR, it was clear that, unlike cells treated with RA alone, cells treated with a combination of RA and TSA re-expressed RAR β mRNA. Under the same experimental conditions, 100 ng/ml TSA alone, or in combination with 1 mM RA, produced 77 and 92% growth inhibition, respectively. Treatment with 1 μ M RA alone did not affect significantly growth inhibition (52%). By MSP analysis, it was assessed that RAR β expression was restored in the presence of a methylated RAR β 2 promoter. This finding indirectly shows that global alterations of HDAC activity, generated by TSA in MDA-MB-231 cells, involved RAR β 2 resulting in RA-induced RAR β expression. Further, demethylation at RAR β 2 did not seem to be an absolute requirement for RAR β gene expression in MDA-MB-231 cells. Noteworthy, persistence of methylation at RAR β 2 was observed also in MCF7 cells where RAR β transcription could be restored in the presence of RA. Growth inhibition was observed in cells treated with TSA alone, or in combination, with RA. Very likely, RAR β along with TSA, a drug known to induce growth inhibition (Yoshida *et al.*, 1995), contributed to the massive growth inhibitory effect that was observed.

[0170] These results show that RAR β 2 promoter is methylated in breast cancer. This study presents evidence that, in breast cancer cells, RAR β 2 promoter undergoes DNA hypermethylation, an epigenetic change known to induce chromatin modifications and influence gene expression. Methylation of the RAR β 2 promoter region was detected, both in breast carcinoma cell lines, and a significant proportion of primary breast tumors. RAR β 2 methylation status did not correlate with the ER status of breast cancer cells and was observed both in *in situ* lesions and invasive tumors. It is not clear when epigenetic changes occur during breast cancer progression. However, methylation of the promoter was not detected in both mortal, and immortal human mammary epithelial cell (HMEC) strains, as well as in normal microdissected breast epithelial cells. These results suggest that aberrant methylation of the RAR β 2 CpG island may be a later event following immortalization. Treatment of breast cancer cells presenting with a methylated RAR β 2, with the demethylating agent 5-Aza-CdR, induced partial DNA demethylation and restored RAR β gene expression. This evidence clearly indicates that DNA methylation is at least a component contributing to RAR β downregulation/loss.

EXAMPLE 5

Hypermethylation of HOXA5

[0171] The extent of methylation of the HOXA5-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors was examined.

[0172] **Tissue preparations and cells** Freshly excised primary breast carcinomas or mammaplasty specimens were minced fine with razor blades and digested with 0.15% collagenase A and 0.5% dispase II (Boehringer Mannheim) prepared in RPMI 1640 medium. The cell clumps were separated from the lighter fibroblasts by gravity separation 3 times. The cell clumps were then digested for 15' with trypsin, washed, and immunostained with anti-cytokeratin-specific antibody (CAM 5.2, Becton-Dickinson) to assess the level of epithelial cell enrichment. The epithelial cells comprised between 70-80% of the enriched cell population.

[0173] Frozen, surgically excised breast tumor samples were cryosectioned, and representative sections were screened by a pathologist after staining with hematoxylin and eosin. Sections containing more than 70% carcinoma cells were used for RNA and protein extractions directly. Breast cancer cell lines and immortalized HMECs were obtained from ATCC (Rockville, MD). Finite life span HMECs were obtained from Dr. Martha Stampfer, HMEC strain 9F1403 was obtained from Clonetics.

[0174] **Methylation specific PCR (MSP) and sodium bisulfite DNA sequencing** One µg of genomic DNA was treated with sodium bisulfite²¹ and was analyzed for MSP using primer sets specific for methylated DNA: 5'-TTTAGCGGTGGCGTTCG-3' (sense; SEQ ID NO:69) and 5'-ATACGACTTCGAATCACGTA-3' (antisense; SEQ ID NO:70), and primers specific for unmethylated DNA: 5'-TTGGTGAAGTTGGGTG-3' (sense; SEQ ID NO:71), and 5'-AATACAACTTCAAATCACATAC-3' (antisense; SEQ ID NO:72) which yielded products of 183 and 213 bp respectively. Sodium bisulfite treated DNA was used to PCR-amplify the HOXA5 promoter region -97 to -303 bp, using the primers

1005333-012023

5'-ATTTTGTATAATGGGTTGTAAT-3' (sense; SEQ ID NO:73) and 5'-AACATATACTTAATTCCCTCC-3' (antisense; SEQ ID NO:74). The product was purified using a Qiagen PCR purification kit (Qiagen Corp.) and was sequenced using the sense primer with an ABI automated fluorescent sequencer according to the manufacturer's instructions.

[0175] Treatment of cells with 5'-aza-2'-deoxycytidine (5-aza-dC) MDA-MB-231 breast cancer cells were treated with 0.75 μ M 5-aza-dC (Sigma), and collected at 0, 3 and 5 days later. RT-PCR was performed using primers: 5'-TCATTTTGCGGTCGCTATCC-3' (sense; SEQ ID NO:75) and 5'-GCCGGCTGGCTGTACCTG-3' (antisense; SEQ ID NO:76).

[0176] Immunoblot Analysis Proteins were visualized by Western analysis and 10% SDS-PAGE. The primary antibodies [anti-HOXA5 (HOXA5-2, BABCO), anti-p53 (AB-6, Oncogene Science), or anti- β -actin (AC-15, Sigma), anti-p21 (15091A, Pharmingen), anti-Mdm2 (65101A, Pharmingen), anti-PARP (AB-2, Oncogene Sciences), anti-dynein (Zymed) and anti- Na⁺, K⁺-ATPase (Ed Benz, Johns Hopkins) (also used as loading controls, with actin)] were used at 1:1000 dilution.

[0177] p53 inactivation by mutation is low (20%) in human breast cancer. Looking for other mechanisms that may account for loss of p53 function in these tumors, the levels of p53 mRNA in breast cancer cell lines and in primary tumors was examined. p53 mRNA levels were 5-10 fold lower in tumor cells than in normal breast epithelium. A subsequent study looked for a consensus protein binding sites in the p53 promoter (Reisman, *et al. Proc. Natl. Acad. of Sci. USA* 85:5146-5150 (1988)), including those of HOX proteins which are known to function as transcription factors (Deschamps, *et al. Crit. Rev. Oncog.* 3:117-173 (1992); Scott, *Nat. Genetics* 15:117-118 (1997)). Selected HOX genes are differentially expressed in neoplasms of a number of tissues, but their functional relationship to the neoplastic phenotype remains to be elucidated. Six putative HOX-core binding sequences (ATTA) were identified within the 2.4 kb human p53 promoter. Of a number of HOX genes examined in breast tumor cells and control breast epithelium, HOXA5 mRNA levels were drastically reduced in breast cancer cells. In fact, there was a tight correlation between p53 and HOXA5

2025 RELEASE UNDER E.O. 14176

mRNA levels in the ten cell lines tested for both genes, with a correlation coefficient $r=0.942$. No such decreased expression was observed for HOXA10, B3, B7, or C8 mRNAs.

[0178] To test for a causal relationship between the decreased expression of p53 and HOXA5 mRNAs, ZR75.1 breast cancer cells or SAOS2 osteosarcoma cells were co-transfected with the -356 bp or the -2.4 kb human p53 promoter-Luciferase reporter together with HOX expression plasmids. HOXA5 transactivated the p53 promoter-dependent reporter activity up to 25-fold in ZR75.1 cells and up to 7-fold in SAOS2 cells. This effect was not seen with other homeotic genes HOXB4, HOXB5 and HOXB7.

[0179] Positive regulation of transcription by HoxA5 was observed with the mouse p53 promoter as well. A single putative Hox-binding sequence (located at nts -204 to -201) was identified in the upstream regulatory region of the murine p53 gene. SAOS2 cells were cotransfected with a -320 bp mouse p53 promoter fused to the CAT gene, together with expression plasmids encoding full-length murine HoxA5, HoxA7, or HoxC8 proteins. Similar to human HOXA5, a 15- to 20-fold increase in CAT activity in the SAOS2 cells cotransfected with the HoxA5 expression plasmid was observed, but no significant effect of HoxA7 or HoxC8. These results suggest that expression from the mouse p53 promoter is specifically stimulated by HoxA5. To define the sequence requirements for the transactivation function, a deletion construct of the p53-promoter CAT construct was tested in cotransfection assays with the full-length HoxA5 expression plasmid. A deletion to -153 bp in the promoter region of the p53-CAT construct eliminated stimulation of CAT activity by the effector plasmid. A truncated HoxA5 protein termed pCMV Δ HoxA5, lacking the homeodomain, was completely inactive in these experiments. Finally a "TT" to "GG" mutation in the core-binding site (-320 mp53MutCAT) that abolished DNA/protein complex formation in cell extracts (see below), completely abrogated transactivation of the CAT reporter gene by HoxA5.

[0180] Direct binding of HoxA5 to the ATTA-containing site in the p53 promoter (positions -204 to -201) was demonstrated by electrophoretic mobility shift (EMSA) and supershift assays. A band was observed in cell extracts from HoxA5 transfected cells, but not in extracts from control cells. This band was competed out by an excess of unlabeled

oligonucleotide but not by an oligonucleotide with an unrelated sequence. No protein/DNA complex was observed in extracts mixed with an oligonucleotide primer which carries two mutations (TT to GG) in the core binding site. Finally, HOXA5 antibodies, but not pre-immune serum, caused a supershift of the bound HOXA5 protein/oligonucleotide complex. This supershift was abrogated by pre-incubation with excess antibody (antigen depletion). Similar shift patterns were observed in extracts of RKO cells transfected with the effector plasmid. These results indicate that the ATTA-containing sequence in the mouse p53 promoter is indeed a HoxA5-binding sequence.

[0181] The above results suggest that HOXA5 may possess growth-suppressive properties through activation of p53 expression. To test this possibility, breast cancer cells, MCF-7 and ZR75.1, which harbor wildtype p53 genes, were transfected with the full length HOXA5 and the Δ HOXA5 (homeodomain-deleted) expression plasmids and tested for colony-forming ability. No surviving colonies were obtained from HOXA5-transfected cells whereas those transfected with Δ HOXA5 and the vector control generated colonies with equal efficiency. To obtain stable cultures that could express HOXA5, clones of MCF-7 cells were generated in which the *HOXA5* gene was placed under the control of an ecdysone-inducible promoter. Within 3 hours after induction of HOXA5 expression by the ecdysone analog, Ponasterone A (Pon A), the levels of p53 mRNA rose by 2-fold. Western blotting showed that p53 and its downstream targets, p21 and Mdm2, as well as HOXA5 were reproducibly induced 2-5 fold following treatment with Pon A. Moreover, addition of Pon A resulted in cell shrinkage by 24 hours followed by significant cell death (80-90%) after 48 hours. Cell death occurred by apoptosis according to the following criteria: 1) cells shrank and formed contractile bodies; 2) DNA laddering was observed; 3) poly (ADP-ribose) polymerase, a substrate for caspases, underwent cleavage by 12 hours; and 4) 70% of the cells showed micronucleus formation, membrane blebbing, and ghost cell features upon staining with acridine orange. This apoptosis was not accompanied by a detectable change in the levels of Bax protein.

[0182] The results herein are consistent with the hypothesis that an increase in the level of HOXA5 in MCF-7 cells leads to an increase in p53 levels, which in turn results in apoptosis. As a further proof of this model, MCF-7 cells expressing the E6 gene of human papilloma

virus, when transfected with the HOXA5 expression vector, were fully able to form colonies. Presumably, the induced p53 in these cells was sequestered by E6 protein and was unable to induce apoptosis. These results support the idea that HOXA5 induces apoptosis through a p53-dependent pathway in MCF-7 cells. This is the first demonstration of the involvement of a HOX protein in apoptosis.

[0183] The hypothesis that HOXA5-induced apoptosis is mediated by p53 was tested as follows. The p53+/+ HCT 116 line of colon carcinoma cells and its p53-/- derivative clone 379.2 were transfected with HOXA5 and p53 expression vectors. Expression of HOXA5 or p53 in the parental HCT116 cells reduced the ability of the cells to form colonies. In contrast, HOXA5 and p53 expression led to different phenotypes in p53 null 379.2 cells. Whereas expression of p53 in these cells abrogated colony formation, expression of HOXA5 had no detectable effect. In the HOXA5-transfected cultures, stable colonies, expressing detectable amounts of HOXA5 protein, and of a size and number comparable to the vector control were observed. Thus, HOXA5 induces cell death only in the presence of a wild-type p53 gene, adding further evidence that p53 mediates HOXA5 activity. Conversely, cells lacking HOXA5 and p53 would be unable to mount a normal response to treatments, such as DNA damage, that normally raise p53 levels by stabilizing the protein. To test this possibility, the two tumor cell lines 21PT and 21MT, which have low expression of HOXA5 and p53 were treated with γ - radiation. No detectable increase in p53 level in 21 PT and 21 MT was observed, while, as expected, p53 was induced in MCF-7 cells.

[0184] These findings in cell culture experiments have *in vivo* correlates. In sixty-seven percent (20/30) of primary breast tumors, HOXA5 protein was undetectable. Strikingly, concurrent loss of p53 expression was observed in the same tumors that lacked HOXA5. Among those tumors expressing HOXA5, one showed a band migrating faster than wild-type HOXA5 present in the RKO cells. HOXA5 cDNAs from eleven p53-negative breast cancer samples and two finite life span human breast epithelial cell (HMEC) strains were sequenced. All HOXA5 coding regions were wild type, except that of tumor #5 which contained a frameshift mutation (G insertion at codon 204) that created a premature stop codon. There is

a coupled loss of p53 and HOXA5 expression in primary breast carcinomas, possibly due to lack of expression or mutational inactivation of HOXA5.

[0185] Seeking an explanation for the absence of the protein in the tumors, HOXA5 DNA of 20 HOXA5-negative and 5 HOXA5-positive primary tumors was sequenced. All contained the wild-type sequence except tumor #5, in which the insertion of G was again found and which contained no wild-type allele. In the absence of mutations, loss of HOXA5 may be a consequence of a loss of upstream regulatory factors or may reflect some repressive phenomenon such as methylation of the gene. Methylation specific PCR (MSP) of sodium bisulfite-treated DNA showed that 16/20 of the tumors contained partially or completely methylated CpGs in the HOXA5 promoter region (ACCN No. AC004080). In contrast, this region was completely unmethylated in human mammary epithelial cells (HMEC) of finite life span, 184 and 9F1403, and in 4 immortalized HMECs, HBL100, MCF10A, 184B5 and 184A1. Nucleotide sequencing of the region -97 bp to -303 bp of the HOXA5 promoter, using sodium bisulfite-treated DNA from HMEC 184, and cancer cell lines, MCF-7 and MDA-MB-231, showed that methylation correlated with silencing of gene expression. Expression of HOXA5 mRNA could be re-initiated in MDA-MB-231 cells by treatment with the DNA methyl transferase inhibitor, 5-aza-2'-deoxycytidine (5-aza-dC). These results are strong preliminary evidence that methylation of the HOXA5 promoter region may be responsible for silencing of gene expression.

[0186] Unlike most tumor types, up to 80% of sporadic breast cancers do not contain p53 mutations. These results suggest that the reduced p53 levels in these tumors result from the absence of a positive regulator of p53 mRNA synthesis. p53 normally functions as a tetramer, so even a small reduction in the concentration of p53 monomers can greatly reduce the effective concentration of tetramers. These results show for the first time that transfected HOXA5 upregulates both p53 promoter-reporter constructs and endogenous p53 synthesis, leading to apoptosis. Finally, HOXA5 was detectable in only one-third of the primary tumors. In the majority of the remaining tumors, lack of HOXA5 expression strongly correlated with methylation of its promoter region, suggesting a causal role for methylation in the silencing of HOXA5 gene expression.

[0187] In summary, these experiments show that HOXA5 is a positive regulator of p53 transcription and function in cultured cells. The correlation observed between HOXA5 and p53 levels in clinical breast cancer demonstrates that loss of HOXA5 expression is an important step in tumorigenesis.

EXAMPLE 6

Hypermethylation of NES-1

[0188] The extent of methylation of the NES-1-associated CpG islands in normal mammary epithelium, in breast cancer cell lines, and in primary mammary tumors was examined.

[0189] **Cell Lines and Tissues** The immortalized HMECs 184A1 (passage 15 and 99) were kindly provided by Dr. Martha Stampfer, and grown as described (<http://www.lbl.gov/LBL-Programs/mrgs/review.html> ;incorporated by reference herein). Mammary organoids were prepared from reduction mammoplasty specimens of women with benign or no abnormalities in the breast following collagenase digestion as described (Bergstraesser and Weitzman (1993) *Cancer Res.*, 53:2644-2654). Primary breast tumor tissues were obtained after surgical resection at the John Hopkins University, and stored frozen at -80°C. DNA was extracted by standard methods (). RNA was extracted with Triazol.

[0190] **Methylation-specific PCR (MSP)** One µg genomic DNA was treated with sodium bisulfite as described in Herman *et al.*(*supra*), and was analyzed by MSP using primer sets located within the third exon of Nes 1 gene. Primers specific for unmethylated DNA were 5'-TTGTAGAGGTGGTGTGTTT-3' (sense; SEQ ID NO:77) and 5'-TTGTAGAGGTGGTGTGTTT-3' (antisense; SEQ ID NO:78) and yielded a 128 base-pairs PCR product. Primers specific for methylated DNA were 5'-TTCGAAGTTTATGGCGTTTC-3' (sense; SEQ ID NO:79) and 5'-TTATTTCCGCAATACGCGAC-3' (antisense; SEQ ID NO:80) and yielded a 137 base-pairs PCR product. The PCR conditions were as follows: 1 cycle of 95°C for 5 min

10035570-012002

[0192] NES-1 expression was observed in mammary organoids and HMEC's from mammaplasty specimens of normal and benign disease breast. In finite life span HMEC primary breast carcinomas analyzed by RT-PCR, NES-1 expression was observed in seven of eleven samples. MSP analysis for a CpG-rich island at NES-1 third exon in the same samples showed methylated sequences in samples that showed NES-1 expression and unmethylated sequences in samples without NES-1 expression. Methylated NES-1 is absent in normal tissue.

68

[0193] In earlier examples use of methylation-specific polymerase chain reaction (PCR) technology (MSP) for detection of the promoter methylation status of human cyclin D2, retinoic acid receptor beta (RAR β), and Twist genes (called “direct MSP”) is described. These genes are essentially unmethylated in normal tissue, but high levels of methylation were found in carcinoma. The present example illustrates a broad study of ductal and lobular carcinoma employing two additional markers, RASSF1A, and Hin-1 genes, in order to achieve the goal of detection of 100% of breast carcinomas. Results of this study show that 100% of invasive ductal carcinoma patients can be detected using the combination of Cyclin D2, RAR β , Twist, and RASSF1A markers (N=27 patient). In addition 100% of invasive lobular carcinoma patients can be detected using the combination of Cyclin D2, RAR β , Twist, and Hin-1 markers (N=19 patients). In the study of 129 patients, the incidence (%) of patients detected with methylation of each of these genes in breast carcinoma is as indicated in Table 3 below.

TABLE 3

Cyc D2	RAR beta	Twist	RASSF1A	Hin1	
19	25	20	62	53	LCIS, in situ
35	20	20	85	79	Lobular carcinoma, invasive
28	33	20	80	75	Grade 1 DCIS, in situ
21	50	23	50	58	Grade 2 DCIS, in situ
42	47	42	78	63	Grade 3 DCIS, in situ
54	30	47	66	59	Ductal carcinoma, invasive

[0194] Thus, the direct MSP technology provides a mechanism for detection of most human breast cancer by molecular methods.

[0195] Potential problems limiting such analyses are mainly the small amount of DNA that is available under certain circumstances (e.g. in ductal lavage, where fluid and cells are obtained from the breast duct) and the need to enhance detection of trace amounts of methylated tumor (e.g. in analyses of blood for circulating tumor DNA, diluted by the presence of a vast excess of unmethylated DNA from blood cells). These problems have now been overcome by development a new technology called multiplex MSP. The procedure for multiplex MSP is basically the following three steps:

1. DNA is isolated and treated with sodium bisulfite, as in direct MSP.
2. PCR reaction #1 is performed using 2 µl DNA (□0.1 µg) in the presence of 5 pairs of primers that will specifically amplify Cyclin D2, RARβ, Twist, RASSF1A, and Hin-1 in the same tube. These primers bind DNA whether or not it is methylated and they bind outside the region that is amplified in PCR reaction #2.

3. PCR reaction #2 is performed using 1 μ l diluted PCR-derived DNA from the first PCR reaction. As in direct MSP, one pair of primers is used per tube that will amplify one gene (either Cyclin D2, RAR β , Twist, RASSF1A, or Hin-1) and the primers are methylation status-specific. Thus two tubes are run per test (patient sample) in PCR reaction #2, each for detection of either unmethylated or methylated DNA respectively. In this reaction PCR-derived DNA is diluted between 10^1 and 10^7 fold (See Figure 11).

[0196] In more detailed terms, multiplex methylation-specific PCR was accomplished by performing two sequential PCR reactions. The first PCR reaction used 5 pairs of gene-specific external primers to co-amplify Cyclin D2, RAR β , Twist, RASSF1A, and Hin-1. The external primer pairs hybridized to sequences outside the region covered by the second PCR reaction. External primers do not contain CpG sequences, thus DNA amplification was independent of methylation status of the genome. The second PCR reaction used 1 pair of gene-specific internal primers to amplify DNA. Unlike the first PCR reaction, for the second PCR reaction primers were methylation status-specific. All primers recognized only sodium bisulfite treated DNA (data not shown). The primer sequences utilized are shown in Table 4 for each gene.

[0197] For the first PCR reaction 2 μ l sodium bisulfite-treated DNA was added to a reaction mixture containing 166 mM (NH₄)₂SO₄, 670 mM Tris, pH 8.8, 67 mM MgCl₂, 100 mM β -mercaptoethanol, 1% DMSO and 4 μ g/ml of each external primer, in a final volume of 25 μ l. The reaction was overlaid with 2 drops oil in a 500 μ l eppendorf tube. Samples were incubated at 95 °C for 5 min, and then 35 cycles of 95 °C for 30 sec, 56 °C for 30 sec, and 72 °C for 45 sec. The final extension was performed at 72 °C for 5 min. For the second PCR reaction, 1 μ l of the first PCR reaction (diluted 1:10²- 1:10⁶) was added to the PCR reaction mix, as described above, which in addition contained 4 μ g/ml of each of two internal primers (forward and reverse). External primers were not added. Reactions to detect methylated and unmethylated genome were carried out in separate reaction tubes, in 8-well strip tubes covered with 2 drops of oil/well. PCR reaction conditions were identical to the first reaction.

[0198] Using this technique, it was determined that multiplex MSP greatly enhances the amount of DNA available for analyses of markers of tumor methylation. The test capacity for direct MSP if $\sim 1 \mu\text{g}$ starting DNA is used enables evaluation of 5 genes in duplicate. By comparison, if $\sim 0.1 \mu\text{g}$ of starting DNA is used in multiplex MSP, a panel of 5 genes can be evaluated in 25 replicate tests, and there is the potential that 10 panels of 5 genes in replicates of 25 tests could be evaluated from $\sim 1 \mu\text{g}$ starting DNA. This would be true if the PCR reaction DNA was conservatively diluted only 10^1 fold, and we have observed that it may be possible to dilute it much higher (i.e. 10^5 - 10^6 fold) to further enhance the availability of sample DNA.

[0199] Multiplex MSP was found to be highly specific, demonstrating concordance with direct MSP analyses of samples obtained from normal human white blood cells (WBC), breast cancer cell lines, and primary breast tumors. Samples found unmethylated by direct MSP were unmethylated by multiplex MSP as well. Furthermore, higher sensitivity for detection of methylated DNA was observed with multiplex MSP, as traces of methylated DNA were detectable by multiplex MSP that were not detectable by direct MSP in some samples.

[0200] In these studies, Cyclin D2, ASSF1A and/or Twist were found to be methylated (at least one marker) in 100% of invasive ductal carcinomas in a sample of 27 cell lines tested. Also gene promotor methylation was found in invasive lobular carcinoma cells as follows: RASSF1A = 85% (n=20); HIN-1 = 79% (n=19); Twist = 20% (n=20); RAR β = 20% (n=20); and CyclinD2 = 35% (n=20). Gene promotor methylation was found in invasive ductal carcinoma cells as follows: RASSF1A = 66% (n=20); HIN-1 = 59% (n=19); Twist = 47% (n=20); RAR β = 30% (n=20); and CyclinD2 = 54% (n=20). The incidence of various combinations of cyclin D2, RAR β , Twist, TASSF1A and HIN-1 in invasive ductal carcinoma in a study of breast cancer cell lines (n=27) was also determined using Multiplex methylation-specific PCR. The combination of cyclin D2, RAR β and Twist occurred in 89% of the samples; the combination of cyclin D2, RAR β , Twist and RASSF1A occurred in 100% of the samples; and the combination of Cyclin D2, RAR β , Twist, and Hin-1 occurred in 93% of the samples tested. The combination of RASSF1A and HIN-1 detected invasive lobular

carcinoma with 95% accuracy. These studies show that RASSF1A and HIN-1 are preferred markers for evaluating a subject having or suspected of having early stage tumorigenesis of breast tissue and that a Multiplex methylation-specific PCR assay utilizing the five markers RASSF1A, Twist and Cyclin D2 will provide an accuracy of 100% detection of invasive ductal carcinoma.

[0201] In conclusion, the multiplex MSP technology can greatly enhance the detection of trace amounts of methylated DNA from patient samples, in a manner which is highly specific. Multiplex MSP can also greatly increase the amount of DNA available for analyses of a wider number of markers of tumor methylation than can presently be analyzed by direct PCR. This technology could allow for analyses of up to 50 genes (10 panels of 5 genes) from the same amount of starting material that can maximally be used to analyze 5 genes using direct MSP.

[0202] Although the invention has been described with reference to the presently preferred embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

RECEIVED 01/15/01

TABLE 4

SEQ ID NO:	Gene	Sense/ antisense		
1	WT	Sense	5'-GCGGGCGCAGTTCCCCAACCA-3'	nucleotides 882-901
2	WT	antisense	5'-ATGGTTTCTCACCAGTGTGCTT-3'	nucleotides 1416-1437
3	WT	Sense	5'-GCATCTGAAACCAGTGAGAA-3'	nucleotides 1320-1339
4	WT	antisense	5'-TTTCTCTGATGCATGTTG-3'	nucleotides 1685-1702
5	WT	Sense	5'-GATTGGCTACCCAAGTGTGCA-3'	
6	WT	antisense	5'-CAGGGGCAGCAGCCACAAAGGC-3'	
7	WT	sense	5'-TTTGGGTAAAGTTAGGCGTCGTCG-3'	
8	WT	antisense	5'-ACACTACTCCTCGTACGACTCCG-3'	
9	WT	sense	5'-TTTGGGTAAAGTTAGGTGTTGTTG-3'	
10	WT	antisense	5'-ACACTACTCCTCATACAACTCCA-3'	
11	WT	sense	5'-CGTCGGGTGAAGGCGGGTAAT-3'	
12	WT	antisense	5'-CGAACCCGAACCTACGAAACC-3'	
13	WT	sense	5'-TGTTGGGTGAAGGTGGGTAAT-3'	
14	WT	antisense	5'-CAAACCCAAACCTACAAAACC-3'	
15	cyclin D2	sense	5'-CATGGAGCTGCTGTGCCACG-3'	
16	cyclin D2	antisense	5'-CCGACCTACCTCCAGCATCC-3'	
17	cyclin D1	sense	5'-AGCCATGGAACACCAGCTC-3'	
18	cyclin D1	antisense	5'-GCACCTCCAGCATCCAGGT-3'	
19	cyclin D2	sense	5'-GATTGGCTAC CCAACTGTTGCA-3'	
20	cyclin D2	antisense	5'-CAGGGGCAGCAGCCACAAAGGC-3'	
21	cyclin D2	sense	5'-GTTATGTTATGTTTGTTGTATG-3'	unmethylated
22	cyclin D2	antisense	5'-GTTATGTTATGTTTGTTGTATG-3'	unmethylated
23	cyclin D2	sense	5'-TACGTGTTAGGGTCGATCG-3'	methylated
24	cyclin D2	antisense	5'-CGAAATATCTACGCTAAACG-3'	methylated
129	cyclin D2	sense	5'-TATTTTTTGTAAAGATAGTTTTGAT-3'	External
130	cyclin D2	antisense	5-TACAACTTTCTAAAAATAACCC-3'	External
25	14.3.3 sigma	sense	5'-ACAGGGGAACTTTATTGAGAGG-3'	A 375 bp σ -specific probe
26	14.3.3 sigma	antisense	5'-AAGGGCTCCGTGGAGAGGG-3'	(SEQ ID NO:26)
27	14.3.3 sigma	sense	5'-GAGGAGTGTCCCGCCTTGTGG-3'	A TG repeat sequence in the 3'UTR of σ
28	14.3.3 sigma	antisense	5'-GTCTCGGTCTTGCACTGGC3'	

SEQUENCE LISTING

SEQ ID NO:	Gene	Sense/ antisense	.	
29	14.3.3 sigma	sense	5'-GTGTGTCCCCAGGCCATGG-3'	A 1.2 kb PCR product, encompassing the entire σ coding sequence, was generated using two primers
30	14.3.3 sigma	antisense	5'-GTCTCGGTCTTGCACTGGCG-3'	(antisense; SEQ ID NO:30)
31	14.3.3 sigma	antisense	5'-CACCTTCTCCCGGTACTCACG-3'	entire σ coding sequence:
32	14.3.3 sigma	sense	5'-GAGCTCTCCTGCGAAGAG-3'	entire σ coding sequence:
33	14.3.3 sigma	sense	5'-GAGGAGGCCATCCTC TCTGGC-3'	entire σ coding sequence:
34	14.3.3 sigma	antisense	5'-TCCACAGTGTGAGGTTGTCTCG-3'	entire σ coding sequence:
35	14.3.3 sigma, first exon	sense	5'-GAGAGAGTTAGTTTGATTTAGAAG-3'	start at nt 8641 generates a 474 bp PCR product
36	14.3.3 sigma	antisense	5'-CTT ACTAATATCCATAACCTCC-3'	(antisense primer with start at nt 9114;
37	14.3.3 sigma	sense	5'-TGGTAGTTTTTATGAAAGGCGTC-3'	methylated DNA
38	14.3.3 sigma	antisense	5'-CCTCTAACCGCCACCACG-3'	
39	14.3.3 sigma	sense	5'-ATGGTAGTTTTTATGAAAGGTGTT-3'	unmethylated DNA
40	14.3.3 sigma	antisense	5'-CCCTCTAACCACCCACCACA-3'	
41	14.3.3 sigma	sense	5'-GTGTGTCCCCAGGCCATGG-3'	PCR was performed using the σ -specific primers

2003-10-06 10:00:00

SEQ ID NO:	Gene	Sense/ antisense		
42	14.3.3 sigma	antisense	5'-ACCTTCTCCCGGTACTCACG-3'	
43	RAR β	sense	5'-AGA GTT TGA TGG AGTTGG GTG GAG-3'	227 bp probe was amplified
44	RAR β	antisense	5'-CAT TCG GTT TGGGTC AAT CCA CTG-3'	
45	RAR β	sense	5'-CAGCCCGGGTAGGGTTCACC-3'	W3
46	RAR β	antisense	5'-CCGGATCCTACCCCGACGG-3'	W3
47	RAR β	sense	5'-CCGAGAACGCGAGCGATCC-3'	W4
48	RAR β	anti-sense	5'-GGCCAATCCAGCCGGGGCG-3'	W4
49	RAR β	sense	5'-GTG GGT GTA GGT GGA ATA TT-3'	unmethylated DNA were as follows: U1
50	RAR β	antisense	5'-AAC AAA CAC ACA AAC CAA CA-3'	U1
51	RAR β	sense	5'-TGT GAG TTA GGA GTA GTG TTTT-3'	U2
52	RAR β	antisense	5'-TTC AAT AAA CCC TAC CCA-3'	U2
53	RAR β	sense	5'-TTA GTA GTT TGG GTA GGGTTT ATT-3'	U3
54	RAR β	antisense	5'-CCA AAT CCT ACC CCAACA-3'	U3
55	RAR β	sense	5'-GAT GTT GAG AAT GTGAGT GAT TT-3'	U4
56	RAR β	antisense	5'-AAC CAA TCC AACCAA AAC A-3'	U4
57	RAR β	sense	5'-AGC GGGCGT AGG CGG AAT ATC-3'	methylated M1
58	RAR β	antisense	5'-CAACGA ACG CAC AAA CCG ACG-3'	M1
59	RAR β RAR β	sense	5'-CGT GAG TTA GGA GTA GCG TTT C-3'	M2
60	RAR β	antisense	5'-CTT TCG ATA AAC CCT ACC CG-3'	M2
61	RAR β	sense	5'-GGT TAG TAG TTC GGG TAG GGT TTA TC-3'	M3
62	RAR β	antisense	5'-CCG AAT CCT ACC CCGACG-3'	M3
63	RAR β	sense	5'-GTC GAG AAC GCG AGCGAT TC-3'	M4
64	RAR β	antisense	5'-CGA CCA ATC CAA CCGAAA CG-3'	M4
65	RAR β	sense	5'-GAC TGT ATG GAT GTTCTG TCA G-3'	RT \pm PCR exon 5
66	RAR β	antisense	5'-ATT TGTCTT GGC AGA CGA AGC A-3'	exon 6
133	RAR β	sense	5'-GTAGGAGGGTTTATTT TTTGTT-3'	External
134	RAR β	antisense	5'-AATTACATTTTCCAAACTTACTC-3'	External
135	RAR β	sense	5'-GGATTGGGATGTTGAGAATGT-3'	Methylated
136	RAR β	antisense	5'-AACCAATCCAACCAAAACAA-3'	Methylated
92	RAR β	sense	5'-GGATTGGGATGTTGAGAATGT-3'	Unmethylated
93	RAR β	antisense	5'-CAACCAATCCAACCAAAACAA-3'	Unmethylated
67	Actin	sense	5'-ACC ATG GAT GAT GAT ATCG-3'	RT \pm PCR
68	Actin	antisense	5'-ACA TGG CTG GGG TGT TGA AG-3'	
69	HOXA5	sense	5'-TTTAGCGGTGGCGTTCG-3'	methylated DNA
70	HOXA5	antisense	5'-ATACGACTTCGAATCACGTA-3'	
71	HOXA5	sense	5'-TTGGTGAAGTTGGGTG-3'	unmethylated

200210162965001

SEQ ID NO:	Gene	Sense/ antisense		
72	HOXA5	antisense	5'-AATACAACCTTCAAATCACATAC-3'	
73	HOXA5	sense	5'-ATTTTGTATAATGGGTTGTAAT3'	
74	HOXA5	antisense	5'-AACATATACTTAATTCCTCC-3'	
75	HOXA5	sense	5'-TCATTTTGC GGTCGCTATCC-3'	RT-PCR
76	HOXA5	antisense	5'-GCCGGCTGGCTGTACCTG-3'	
77	NES-1	sense	5'-TTGTAGAGGTGGTGTGTTT-3'	unmethylated
78	NES-1	antisense	5'-CACACAATAAAACAAAAACCA -3'	
79	NES-1	sense	5'-TTCGAAGTTTATGGCGTTTC-3'	Methylated
80	NES-1	antisense	5'-TTATTTCCGCAATACGCGAC-3'	
81	NES-1	sense	5'-ACCAGAGTTGGGTGCTGAC-3'	
82	NES-1	antisense	5'-ACCTGGCACTGGTCTCCG-3'	
83	36B4	sense	5'-GATTGGCTACCCAACCTGTTGCA-3'	
84	36B4	antisense	5'-CAGGGGCAGCAGCCACAAAGGC-3'	
85	Estrogen Receptor	sense	5'-G GGTGTTTTTG AGATTGTTGG-3	Unmethylated
86			5'-TG AGTTGTGATG GGTTTTGG-3	
87		antisense	5'-CCAAAACC CATCAAACT CA-3	
88		sense	5'-AGAGTAGGCG GCGAGCGT-3	Methylated
89			5'-CGGGAAAAG TACGTGTTTCG T-3	
90		antisense	5'-A CGAACACGTA CTTTTCCCG-3	
107	Twist	sense	5'-T TTCGGATGGG GTTGTTTCATC-3	Methylated
108	Twist	antisense	5'-AAACGAC CTAACCCGAA CG-3	Methylated
109	Twist	sense	5'-TT TGGATGGGGT TGTTATTGT-3	Unmethylated
110	Twist	antisense	5'-C CTAACCCAAA CAACCAACC-3	Unmethylated
133	Twist	sense	5'-GAGATGAGATATTATTTATTGTG-3	External
134	Twist	antisense	5'-AACAACAATATCATTAACCTAAC-3	External
111	HIN-1	sense	5'-AGGGAAGTTTTTTTATTGTTT-3	
112	HIN-1	antisense	5'-GTGGTTTTGTTTGTATGTTTTGGTG-3	
113	HIN-1	antisense	5'-CACCGAAACATACAAAACAAACCAC-3	
114	HIN-1	sense	5'-GTTTGTTAAGAGGAAGTTTT-3	External
115	HIN-1	antisense	5'-CACCGAAACATACAAAACAAACCAC-3	External
116	HIN-1	sense	5'-GGTACGGGTTTTTACGGTTCGTC-3	Methylated
117	HIN-1	antisense	5'-AACTTCTTATACCCGATCCTCG-3	Methylated
118	HIN-1	sense	5'-GGTATGGGTTTTTATGGTTTGTT-3	Unmethylated
119	HIN-1	antisense	5'-CAAACTTCTTATACCCAATCCTCA-3	Unmethylated
122	RASSF1A	sense	5'-GGGAGTTTGAGTTTATTGAGT-3	External
123	RASSF1A	antisense	5'-ACCCCTTAACCTACCCCTTC-3	External
124	RASSF1A	sense	5'-GTTGGTATTC-3	Methylated
125	RASSF1A	sense	5'-GTTGGGCGC-3	Methylated
126	RASSF1A	antisense	5'-GCACCACGTATACGTAACG-3	Methylated
127	RASSF1A	sense	5'-GGTTGTATTGGTTGGAGTG-3	Unmethylated
128	RASSF1A	antisense	5'-CTACAAACCTTACACACAACA-3	Unmethylated

100557-0300

TABLE 6

Multiplex Is Highly Specific

•Concordance Observed Between Direct PCR and Multiplex PCR in
Human WBC DNA Analyses

WBC	Cyclin D2		RARbeta		Twist		RASSF1A		Hin-1	
	Direct	Multi Diln	Direct	Multi Diln	Direct	Multi Diln	Direct	Multi Diln	Direct	Multi Diln
7157	U	5	U	6	U	6	U	6	U	5
7160	U	3	U	5	U	2	U	5	U	3
7163	U	5	U	6	U	6	U	6	U	5
7164	U	5	U	6	U	6	U	6	U	5
H20	NR	1	NR	1	NR	0	NR	1	NR	1

TABLE 7

Multiplex Is Highly Sensitive

•Methylated Signals Not Observed by Direct PCR are Revealed
by Multiplex PCR in Human Breast CA Cell Line Analyses

Cell Lines	Cyclin D2			RARbeta			Twist		RASSF1A			Hin-1		Multi		Diln
	Direct	Multi	Diln	Direct	Multi	Diln	Direct	Multi	Direct	Multi	Diln	Direct	Multi	Diln	Diln	
7160	U	U	4	U	U	4	U	U/M	U	U	4	U	U	U	4	4
231	M	M	4	M	M	4	M	M	M	M	4	M	M	M	M	4
MCF-7	U/M	U/M	4	M	M	4	M/Uw	M	M	M	4	Mw	M	M	M	4
MCF-10A	M	M/Uw	4	U/M	U/M	4	U	U/Mw	M	M	4	U/M	U/M	U/M	M	4
HBL100	M	U/M	4	U/Mw	U/M	4	U	U/Mw	M	U/M	4	M/Uw	M	M	M	4
ZR75-1	U	U/Mw	4	M	M/Uw	4	U/M	U/M	M	M	4	M	M	M	M	4